

ON GROWTH CHARACTERISTICS OF OPERATOR-VALUED FUNCTIONS

S.N. MISHIN

Abstract. In the work we generalize Liouville theorem and the concept of order and type of entire function to the case of an operator-valued function with values in the space $\text{Lec}(\mathbf{H}_1, \mathbf{H})$ of all linear continuous operators acting from a locally convex space \mathbf{H}_1 to a locally convex space \mathbf{H} with an equicontinuous bornology. We find the formulae expressing the order and type of an operator-valued function in terms of the characteristics for the sequence of the coefficients. Some properties of the order and type of an operator-valued function are established.

Keywords: locally convex space, order and type of sequence of operators, order and type of entire function, equicontinuous bornology, convergence by bornology, operator-valued function.

INTRODUCTION

It is known [3, 4] that if an entire scalar function $f(z) = \sum_{n=0}^{\infty} a_n z^n$ is not a polynomial, the maximum of its modulus $M_f(r) = \max_{|z| \leq r} |f(z)|$ grows faster than any positive power of r as $r \rightarrow \infty$ (Liouville theorem). To estimate the growth of such functions, one usually uses the characteristics (order and type),

$$\rho = \overline{\lim}_{r \rightarrow \infty} \frac{\ln \ln M_f(r)}{\ln r}, \quad \sigma = \overline{\lim}_{r \rightarrow \infty} \frac{\ln M_f(r)}{r^\rho}. \quad (1)$$

At that, the formulae expressing these characteristics in terms of the coefficients

$$\rho = \overline{\lim}_{n \rightarrow \infty} \frac{n \ln n}{-\ln |a_n|}, \quad (\rho e \sigma)^{\frac{1}{\rho}} = \overline{\lim}_{n \rightarrow \infty} n^{\frac{1}{\rho}} \sqrt[n]{|a_n|} \quad (2)$$

are known. This work is devoted to the generalization of these formulae and the Liouville theorem for the case of an entire operator-valued function $F(t) = \sum_{n=0}^{\infty} A_n t^n$ with the values in the space $\text{Lec}(\mathbf{H}_1, \mathbf{H})$ of all linear continuous operators acting from a locally convex space \mathbf{H}_1 into a locally convex space \mathbf{H} . The spaces \mathbf{H}_1 and \mathbf{H} are in general not normable.

1. ENTIRE OPERATOR-VALUED FUNCTIONS AND ANALOGUE TO LIOUVILLE THEOREM

\mathbf{H}_1 and \mathbf{H} are separable locally convex spaces over the field of complex numbers with the topologies defined respectively by the multinorms $\{\|\cdot\|_q\}$, $q \in \mathcal{Q}$ and $\{\|\cdot\|_p\}$, $p \in \mathcal{P}$. Without loss of generality one can regard the multinorms in \mathbf{H}_1 and \mathbf{H} as majorant [2]. By $\mathcal{A} = \{A_n\}_{n=0}^{\infty}$ we denote a sequence of linear continuous operators acting from the locally convex space \mathbf{H}_1

S.N. MISHIN, ON GROWTH CHARACTERISTICS OF OPERATOR-VALUED FUNCTIONS.

© MISHIN S.N. 2013.

Submitted August 16, 2012.

into the locally convex space \mathbf{H} . The sequence \mathcal{A} is called as having an order [1, 5], if there exists a sequence of positive numbers $\{c_n\}_{n=0}^\infty$ such that

$$\forall p \in \mathcal{P} \exists C_p > 0 \exists q(p) \in \mathcal{Q} \forall x \in \mathbf{H}_1 \forall n \in \mathbb{N} : \|c_n A_n(x)\|_p \leq C_p \|x\|_q', \quad (3)$$

i.e., the family of the operators $\{c_n A_n\}$ is equicontinuous.

Let

$$\theta_{\mathcal{A}}(p, q, n) = \sup_{\|x\|_q' \neq 0} \left\{ \frac{\|A_n(x)\|_p}{\|x\|_q'} \right\}, \quad n = 0, 1, 2, \dots$$

(the case $\theta_{\mathcal{A}}(p, q, n) = +\infty$ is not excluded). We denote

$$\beta_{p,q}(\mathcal{A}) = \overline{\lim}_{n \rightarrow \infty} \frac{\ln \theta_{\mathcal{A}}(p, q, n)}{n \ln n}.$$

Definition 1. The number $\beta_p(\mathcal{A}) = \inf_{q \in \mathcal{Q}} \beta_{p,q}(\mathcal{A})$, ($p \in \mathcal{P}$) is called a p -order of the sequence of the operators \mathcal{A} , and the number $\beta(\mathcal{A}) = \sup_{p \in \mathcal{P}} \{\beta_p(\mathcal{A})\}$ is called its order.

If $\beta(\mathcal{A}) = \pm\infty$ and at that the sequence \mathcal{A} has an order, then it is called a sequence of an infinite order.

Remark. Let us note that there is an essential difference between the sequences having an order $\beta(\mathcal{A}) = +\infty$, and that having no order (despite formally $\beta(\mathcal{A}) = +\infty$). If $\beta(\mathcal{A}) = +\infty$, but the sequence $\mathcal{A} = \{A_n\}$ has an order, it is possible to select a sequence of positive numbers $\{c_n\}$ such that condition (3) holds. And one can not select such a sequence for the sequences having no order.

If a sequence of operators \mathcal{A} has a p -order $\beta_p(\mathcal{A}) \neq \pm\infty$, one introduces for it a finer characteristics. Denote

$$\alpha_{p,q}(\mathcal{A}) = \overline{\lim}_{n \rightarrow \infty} n^{-\beta_p(\mathcal{A})} \sqrt[n]{\theta_{\mathcal{A}}(p, q, n)}.$$

Definition 2. The number $\alpha_p(\mathcal{A}) = \inf_{q \in \mathcal{Q}} \alpha_{p,q}(\mathcal{A})$, ($p \in \mathcal{P}$) is called a p -type of a sequence of operators \mathcal{A} at the p -order $\beta_p(\mathcal{A})$.

It is obvious that $\beta_p(\mathcal{A}) \leq \beta(\mathcal{A})$, $\forall p$. It is possible to show [7] that the case when the identity $\beta_p(\mathcal{A}) = \beta(\mathcal{A})$ is valid not for all p , but just for some p , is reduced to the case $\beta_p(\mathcal{A}) = \beta(\mathcal{A})$, $\forall p$ by replacing the multinorm to an equivalent one. This replacement changes neither the order no the type of a sequence of operators. This is why (without loss of generality) we shall consider two cases, either $\beta_p(\mathcal{A}) = \beta(\mathcal{A})$, $\forall p$, or $\beta_p(\mathcal{A}) < \beta(\mathcal{A})$, $\forall p$.

Definition 3. Let a sequence of operators \mathcal{A} has the p -orders $\beta_p(\mathcal{A})$ and the order $\beta(\mathcal{A}) \neq \pm\infty$. The number

$$\alpha(\mathcal{A}) = \begin{cases} \sup_{p \in \mathcal{P}} \{\alpha_p(\mathcal{A})\} & , \quad \beta_p(\mathcal{A}) = \beta(\mathcal{A}), \quad \forall p \\ 0 & , \quad \beta_p(\mathcal{A}) < \beta(\mathcal{A}), \quad \forall p \end{cases}$$

is called a type of the sequence of operators \mathcal{A} at the order $\beta(\mathcal{A})$.

A sequence of operators \mathcal{A} is called belonging to the class $\mathfrak{L}_{\mathbf{H}_1, \mathbf{H}}[b, a]$, (cf. [1, 5]) if its order is less than b or equal to b , but then the type does not exceed a .

Let \mathbf{H} be a complete space. It is known [8] that in this case the space $\text{Lec}(\mathbf{H}_1, \mathbf{H})$ of linear continuous operators acting from \mathbf{H}_1 into \mathbf{H} equipped with an equicontinuous bornology is a complete bornological vector convex space.

Definition 4. An operator-valued function $F : \mathbb{C} \rightarrow \text{Lec}(\mathbf{H}_1, \mathbf{H})$ is called differentiable at a point $t_0 \in \mathbb{C}$ if there exists a limit (w.r.t. the bornology of the space $\text{Lec}(\mathbf{H}_1, \mathbf{H})$)

$$\lim_{t \rightarrow t_0} \frac{F(t) - F(t_0)}{t - t_0}. \quad (4)$$

This limit is called a derivative of the operator-valued function F at the point t_0 and is indicated by $F'(t_0)$.

Definition 5. An operator-valued function $F : \mathbb{C} \rightarrow \text{Lec}(\mathbf{H}_1, \mathbf{H})$ is called entire if its defined and differentiable at each point $t \in \mathbb{C}$.

An entire operator-valued function is obviously continuous everywhere (w.r.t. the bornology of the space $\text{Lec}(\mathbf{H}_1, \mathbf{H})$).

Let

$$\theta_F(p, q, t) = \sup_{\|x\|_q \neq 0} \left\{ \frac{\|F(t)(x)\|_p}{\|x\|_q} \right\}, \quad t \in \mathbb{C}$$

(the case $\theta_F(p, q, t) = +\infty$ is not excluded).

Theorem 1. An entire operator-valued function $F(t)$ is bounded on each closed disk, i.e., the family of the operators $\{F(t)\}_{|t| \leq r}$ is equicontinuous for each $r > 0$.

Proof. We fix an arbitrary $r > 0$. Suppose the function $F(t)$ is entire, and the family $\{F(t)\}_{|t| \leq r}$ is not equicontinuous, i.e., there exists $p_0 \in \mathcal{P}$ such that for each $C > 0$ and for each $q \in \mathcal{Q}$ there exists $t_C = t_C(q)$ such that $|t_C| \leq r$ and $\theta_F(p_0, q, t_C) > C$. We fix an arbitrary $q \in \mathcal{Q}$ and take $C = n$, $n \in \mathbb{N}$. We obtain then a sequence of complex numbers $t_n = t_n(q)$ lying within the disk $|t| \leq r$. At that,

$$\theta_F(p_0, q, t_n) > n, \quad \forall n. \quad (5)$$

By the boundedness of the sequence $\{t_n\}$ there exists a converging subsequence $\{t_{n_k}\}$. It follows from (5) that $\theta_F(p_0, q, t_{n_k}) > n_k$, $\forall k$, i.e., the sequence $\{F(t_{n_k})\}$ is not equicontinuous and thus diverges. But by the continuity of the function F it must converges. We obtain the contradiction. \square

If the function $F(t)$ is entire, then for each fixed $x \in \mathbf{H}_1$, $F(t)(x)$ is an entire function with values in \mathbf{H} . Such function is represented as a power series [9]

$$F(t)(x) = \sum_{n=0}^{\infty} x_n t^n, \quad x \in \mathbf{H}_1, \quad \{x_n\} \subset \mathbf{H}$$

(the sequence $\{x_n\}$ depends on x). We let

$$M_F(p, q, r) = \sup_{|t| \leq r} \theta_F(p, q, t).$$

We define a sequence of operators $A_n : \mathbf{H}_1 \rightarrow \mathbf{H}$ as follows, $A_n(x) = x_n$, $\forall x \in \mathbf{H}_1$. We obtain the expansion of the function $F(t)$ as a power series

$$F(t) = \sum_{n=0}^{\infty} A_n t^n. \quad (6)$$

At that, series (6) everywhere pointwise converges to the function $F(t)$ (for each fixed $x \in \mathbf{H}_1$ the series $\sum_{n=0}^{\infty} A_n(x) t^n$ converges to the function $F(t)(x)$ everywhere). Let us show that $\{A_n\} \subset \text{Lec}(\mathbf{H}_1, \mathbf{H})$ and series (6) converges everywhere to the function $F(t)$ w.r.t. the bornology. First we prove the following theorem.

Theorem 2 (Analogue of Cauchy inequality). The inequality

$$\theta_{\mathcal{A}}(p, q, n) \leq \frac{M_F(p, q, r)}{r^n}, \quad \forall p \quad \forall q \quad \forall n \quad \forall r > 0 \quad (7)$$

holds true.

Proof. Let $p \in \mathcal{P}$, $q \in \mathcal{Q}$, $r > 0$. If $M_F(p, q, r) = \infty$, then inequality (7) holds true. Let $M_F(p, q, r) < \infty$. Since for each fixed x the vector-function $F(t)(x) = \sum_{n=0}^{\infty} A_n(x)t^n$ is entire, then (see, for instance, [9])

$$A_n(x) = \frac{1}{2\pi i} \int_{|\xi|=r} \frac{F(\xi)(x)d\xi}{\xi^{n+1}}, \quad n \in \mathbb{N}.$$

Hence, $\forall p \in \mathcal{P} \forall x \in \mathbf{H}_1 \forall r > 0 \forall n \in \mathbb{N}$ we have

$$\|A_n(x)\|_p \leq \frac{\sup_{|\xi| \leq r} \|F(\xi)(x)\|_p}{r^n} \leq \frac{\sup_{|\xi| \leq r} \theta_F(p, q, \xi)}{r^n} \|x\|'_q = \frac{M_F(p, q, r)}{r^n} \|x\|'_q$$

that yields inequality (7). \square

Since the function $F(t)$ is entire, by Theorem 1 for each $r > 0$ the family $\{F(t)\}_{|t| \leq r}$ is equicontinuous, i.e.,

$$\forall p \in \mathcal{P} \exists C_p > 0 \exists q_p \in \mathcal{Q} \forall x \in \mathbf{H}_1 \forall t \leq r \Rightarrow \|F(t)(x)\|_p \leq C_p \|x\|'_{q_p}.$$

For each p we choose $q_0 = q_0(p)$ such that $\|x\|'_{q_0} \geq \|x\|'_{q_p}$, $\forall x \in \mathbf{H}_1$ (it is always possible since the multinorm is majorant). Then

$$\theta_F(p, q_0, t) = \sup_{\|x\|'_{q_0} \neq 0} \left\{ \frac{\|F(t)(x)\|_p}{\|x\|'_{q_0}} \right\} \leq \sup_{\|x\|'_{q_0} \neq 0} \left\{ \frac{C_p \|x\|'_{q_p}}{\|x\|'_{q_0}} \right\} = \tilde{C}_p(q_0), \quad |t| \leq r.$$

Thus, for each $r > 0$ and each $p \in \mathcal{P}$ there exists $q_0 \in \mathcal{Q}$ such that $\theta_F(p, q_0, t)$ (as functions of t) are bounded in the disk $|t| \leq r$. And it means that

$$\forall r \forall p \exists q_0(p, r) : M_F(p, q_0, r) < \infty.$$

Hence, by Theorem 2,

$$\overline{\lim}_{n \rightarrow \infty} \sqrt[n]{\theta_{\mathcal{A}}(p, q_0, n)} \leq \frac{1}{r}, \quad r > 0. \quad (8)$$

It follows from (8) that either $\beta_p(\mathcal{A}) < 0$ or $\beta_p(\mathcal{A}) = 0$, but then by the arbitrariness of r

$$\alpha_p(\mathcal{A}) = \inf_{q \in \mathcal{Q}} \overline{\lim}_{n \rightarrow \infty} \sqrt[n]{\theta_{\mathcal{A}}(p, q, n)} = 0.$$

Thus, the sequence $\{A_n\}$ belongs to the class $\mathfrak{L}_{\mathbf{H}_1, \mathbf{H}}[0, 0]$ and therefore series (6) everywhere converges to the function $F(t)$ w.r.t. bornology (see [1, 5]).

Theorem 3 (Analogue of Liouville theorem). *Suppose function (6) is entire and satisfies the condition*

$$\exists k \forall p \exists K_p > 0 \exists q(p) \forall r > 0 : M_F(p, q, r) \leq K_p r^k. \quad (9)$$

Then F is an operator-valued polynomial of degree at most k , i.e.,

$$F(t) = \sum_{n=0}^{[k]} A_n t^n.$$

Proof. By inequalities (7), (9) and the definition of the numbers $\theta_{\mathcal{A}}(p, q, n)$ we have

$$\|A_n(x)\|_p \leq \theta_{\mathcal{A}}(p, q, n) \|x\|'_q \leq K_p r^{k-n} \|x\|'_q, \quad \forall p \forall x \in \mathbf{H}_1 \forall r > 0 \forall n, \quad q = q(p).$$

By the arbitrariness of r ,

$$\|A_n(x)\|_p = 0, \quad \forall n > k \forall p \forall x \in \mathbf{H}_1,$$

thus, $A_n = 0$, $\forall n > k$. \square

Theorem 3 shows that if F is an entire transcendental function, then the quantities $M_F(p, q, r)$ grows faster than any positive power as $r \rightarrow \infty$.

2. GROWTH CHARACTERISTICS FOR ENTIRE FUNCTION AND FORMULAE FOR THEIR CALCULATION

Definition 6. Let $F : \mathbb{C} \rightarrow \text{Lec}(\mathbf{H}_1, \mathbf{H})$ be an entire transcendental function. The number $\rho_p(F) = \inf_{q \in \mathcal{Q}} \rho_{p,q}(F)$, where

$$\rho_{p,q}(F) = \overline{\lim}_{r \rightarrow \infty} \frac{\ln \ln M_F(p, q, r)}{\ln r}$$

will be called a p -order of the function F , and the number $\rho(F) = \sup_{p \in \mathcal{P}} \{\rho_p(F)\}$ will be called its order.

If $0 < \rho_p(F) < \infty$, the number $\sigma_p(F) = \inf_{q \in \mathcal{Q}} \sigma_{p,q}(F)$, where

$$\sigma_{p,q}(F) = \overline{\lim}_{r \rightarrow \infty} \frac{\ln M_F(p, q, r)}{r^{\rho_p(F)}},$$

will be called a p -type of the function f at p -order $\rho(F)$.

It can be shown that the case when for some p , $\rho_p(F) < \rho(F)$, while for other $\rho_p(F) = \rho(F)$, is reduced to the case $\rho_p(F) = \rho(F)$, $\forall p$ by the replacement of the multinorm to an equivalent one. This is why (without loss of generality) we shall consider two cases, either $\rho_p(F) < \rho(F)$, $\forall p$, or $\rho_p(F) = \rho(F)$, $\forall p$.

Definition 7. Suppose a function $F(t)$ has p -orders $\rho_p(F)$ and order $0 < \rho(F) < \infty$. The number

$$\sigma(F) = \begin{cases} 0 & , \quad \rho_p(F) < \rho(F), \quad \forall p \\ \sup_{p \in \mathcal{P}} \{\sigma_p(F)\} & , \quad \rho_p(F) = \rho(F), \quad \forall p \end{cases}$$

will be called a type of the function f at the order $\rho(F)$.

Lemma 1. Suppose

$$\forall p \exists q_p \exists a_p, b_p > 0 \exists r_0(p) \forall r > r_0 : M_F(p, q_p, r) < e^{a_p r^{b_p}}. \quad (10)$$

Then

$$\forall p \exists n_0(p) \forall n > n_0 : \sqrt[n]{\theta_A(p, q_p, n)} < \left(\frac{a_p b_p e}{n} \right)^{\frac{1}{b_p}}. \quad (11)$$

Proof. Suppose inequality (10) holds true, then by (7) we have

$$\theta_A(p, q_p, n) < \frac{e^{a_p r^{b_p}}}{r^n}; \quad \forall p \forall r > r_0(p) \forall n. \quad (12)$$

We denote $\mu_p(r) = e^{a_p r^{b_p}} r^{-n}$. It is obvious that

$$\forall p : \mu_p(0) = \mu_p(+\infty) = +\infty.$$

Let us find $\min_{r>0} \{\mu_p(r)\}$,

$$\mu'_p(r) = \mu_p(r) \ln' \mu_p(r)$$

$$\mu'_p(r) = \mu_p(r) (a_p r^{b_p} - n \ln r)'$$

$$\mu'_p(r) = \mu_p(r) \left(a_p b_p r^{b_p-1} - \frac{n}{r} \right)$$

$\mu'_p(r) = 0$ as $r = r_1 = \left(\frac{n}{a_p b_p} \right)^{\frac{1}{b_p}}$. Substituting r_1 in inequality (12), we obtain (11). □

Lemma 2. *Suppose*

$$\forall p \exists q_p \exists a_p, b_p > 0 \exists n_0(p) \forall n > n_0 : \sqrt[n]{\theta_{\mathcal{A}}(p, q_p, n)} < \left(\frac{a_p b_p e}{n} \right)^{\frac{1}{b_p}}. \quad (13)$$

Then

$$\forall p \forall \varepsilon > 0 \exists r_0(p, \varepsilon) \forall r > r_0 : M_F(p, q_p, r) < e^{(a_p + \varepsilon)r^{b_p}}. \quad (14)$$

Proof. By condition (13) $\mathcal{A} \in \mathfrak{L}_{\mathbf{H}_1, \mathbf{H}}[0, 0]$, thus, F is an entire operator-valued function. Let us fix an arbitrary p (and fix by this depending on it q_p, a_p, b_p) and consider the inequality

$$\theta_{\mathcal{A}}(p, q_p, n)r^n < \left(\left(\frac{a_p b_p e}{n} \right)^{\frac{1}{b_p}} r \right)^n.$$

For sufficiently large n

$$\left(\frac{a_p b_p e}{n} \right)^{\frac{1}{b_p}} r < \frac{1}{2}. \quad (15)$$

By $N_p(r)$ we denote the lowest of natural numbers n for which inequality (15) holds true. Let us find the dependence of $N_p(r)$ on r . We have

$$2r \left(\frac{a_p b_p e}{n} \right)^{\frac{1}{b_p}} < 1, \text{ as } n > (2r)^{b_p} (a_p b_p e).$$

Therefore, we can let $N_p(r) = [(2r)^{b_p} (a_p b_p e)] + 1$.

Further, for each fixed $p \in \mathcal{P}$, $t \in \mathbb{C}$ and $x \in \mathbf{H}_1$ we have

$$\|F(t)(x)\|_p \leq \sum_{n=0}^{\infty} \|A_n(x)\|_p |t|^n \leq \sum_{n=0}^{\infty} \theta_{\mathcal{A}}(p, q_p, n) |t|^n \|x\|'_{q_p},$$

hence,

$$\theta_F(p, q_p, t) \leq \sum_{n=0}^{\infty} \theta_{\mathcal{A}}(p, q_p, n) |t|^n,$$

i.e.,

$$\forall p \forall r > 0 : M_F(p, q_p, r) \leq \sum_{n=0}^{\infty} \theta_{\mathcal{A}}(p, q_p, n) r^n = \sum_{n=0}^{N_p(r)-1} \theta_{\mathcal{A}}(p, q_p, n) r^n + \sum_{n=N_p(r)}^{\infty} \theta_{\mathcal{A}}(p, q_p, n) r^n.$$

For $n \geq N_p(r)$ the inequality $\theta_{\mathcal{A}}(p, q_p, n)r^n < \left(\frac{1}{2}\right)^n$ holds true and hence

$$\sum_{n=N_p(r)}^{\infty} \theta_{\mathcal{A}}(p, q_p, n) r^n < \sum_{n=N_p(r)}^{\infty} \left(\frac{1}{2}\right)^n < \sum_{n=0}^{\infty} \left(\frac{1}{2}\right)^n = 2.$$

Since for each fixed p and r

$$\lim_{n \rightarrow \infty} \theta_{\mathcal{A}}(p, q_p, n) r^n = 0,$$

the sequence $\{\theta_{\mathcal{A}}(p, q_p, n)r^n\}$ has a maximal term. Let

$$m_p(r) = \max_{n \geq 0} \{\theta_{\mathcal{A}}(p, q_p, n)r^n\},$$

then

$$\sum_{n=0}^{N_p(r)-1} \theta_{\mathcal{A}}(p, q_p, n) r^n \leq m_p(r) N_p(r).$$

Let us estimate $m_p(r)$. Let $\theta_{\mathcal{A}}(p, q_p, s)r^s$ be a maximal term. Under an unbound increasing of r the index s of maximal term increases unboundedly as well, i.e., $s \rightarrow \infty$ as $r \rightarrow \infty$. If r is sufficiently large, then $s > n_0$, where n_0 is a number in (13).

This is why

$$m_p(r) = \theta_{\mathcal{A}}(p, q_p, s)r^s < \left(\frac{a_p b_p e}{s}\right)^{\frac{s}{b_p}} r^s \leq \max_{\xi \geq 0} \left\{ \left(\frac{a_p b_p e}{\xi}\right)^{\frac{\xi}{b_p}} r^\xi \right\}.$$

We denote

$$\nu_p(\xi) = \left(\frac{a_p b_p e}{\xi}\right)^{\frac{\xi}{b_p}} r^\xi.$$

Clearly,

$$\forall p: \nu_p(0) = 1, \nu_p(+\infty) = 0.$$

Let us find $\max_{\xi \geq 0} \{\nu_p(\xi)\}$. We have

$$\nu'_p(\xi) = \nu_p(\xi) \left(\frac{\ln(a_p b_p e)}{b_p} - \frac{\ln \xi}{b_p} - \frac{1}{b_p} + \ln r \right).$$

$\nu'_p(\xi) = 0$ as $\xi = \xi_1 = (a_p b_p) r^{b_p}$.

$$\nu_p(\xi_1) = e^{a_p r^{b_p}}.$$

Therefore (for sufficiently large r), $m_p(r) < e^{a_p r^{b_p}}$.

Thus,

$$M_F(p, q_p, r) \leq N_p(r)m_p(r) + 2 \leq ((2r)^{b_p}(a_p b_p e) + 1)e^{a_p r^{b_p}} + 2 < e^{(a_p + \varepsilon)r^{b_p}}.$$

□

Theorem 4. *The growth characteristics of function (6) are calculated by the formulae*

$$\rho_p(F) = -\frac{1}{\beta_p(\mathcal{A})}, \quad \forall p, \quad (16)$$

$$\sigma_p(F) = -\frac{\beta_p(\mathcal{A})}{e} (\alpha_p(\mathcal{A}))^{-\frac{1}{\beta_p(\mathcal{A})}}, \quad \forall p, \quad (17)$$

$$\rho(F) = -\frac{1}{\beta(\mathcal{A})}, \quad (18)$$

$$\sigma(F) = \begin{cases} 0 & , \quad \beta_p(\mathcal{A}) < \beta(\mathcal{A}), \quad \forall p \\ -\frac{\beta(\mathcal{A})}{e} (\alpha(\mathcal{A}))^{-\frac{1}{\beta(\mathcal{A})}} & , \quad \beta_p(\mathcal{A}) = \beta(\mathcal{A}), \quad \forall p. \end{cases} \quad (19)$$

Proof. We fix an arbitrary p . Suppose the p -order of the function F equals $\rho_p(F)$. Then

$$\forall p \quad \forall \varepsilon > 0 \quad \exists q_p(\varepsilon) \quad \exists r_0(p, \varepsilon) \quad \forall r > r_0: \quad M_F(p, q_p, r) \leq \exp \{r^{\rho_p(F) + \varepsilon}\}.$$

By Lemma 1 ($b_p = \rho_p(F) + \varepsilon$, $a_p = 1$)

$$\sqrt[n]{\theta_{\mathcal{A}}(p, q_p, n)} < \left(\frac{(\rho_p(F) + \varepsilon) e}{n} \right)^{\frac{1}{\rho_p(F) + \varepsilon}}, \quad \forall n > n_0.$$

By this we successively find

$$\frac{1}{n} \ln \theta_{\mathcal{A}}(p, q_p, n) < \left(\frac{1}{\rho_p(F) + \varepsilon} \right) \ln \left((\rho_p(F) + \varepsilon) e \right) - \frac{\ln n}{\rho_p(F) + \varepsilon} = C_p(\varepsilon) - \frac{\ln n}{\rho_p(F) + \varepsilon},$$

$$\ln \theta_{\mathcal{A}}(p, q_p, n) < C_p(\varepsilon)n - \frac{n \ln n}{\rho_p(F) + \varepsilon},$$

$$\ln \frac{1}{\theta_{\mathcal{A}}(p, q_p, n)} > \frac{n \ln n}{\rho_p(F) + \varepsilon} - C_p(\varepsilon)n = n \ln n \left(\frac{1}{\rho_p(F) + \varepsilon} - \frac{C_p(\varepsilon)}{\ln n} \right), \quad \forall n > n_0. \quad (20)$$

As $n \rightarrow \infty$, the expression in parentheses in (20) tends to $\frac{1}{\rho_p(F)+\varepsilon}$, and for large n

$$\ln \frac{1}{\theta_{\mathcal{A}}(p, q_p, n)} > \frac{n \ln n}{\rho_p(F) + 2\varepsilon},$$

i.e.,

$$\rho_p(F) + 2\varepsilon > \frac{n \ln n}{-\ln \theta_{\mathcal{A}}(p, q_p, n)}.$$

By the arbitrariness of ε ,

$$-\frac{1}{\beta_{p, q_p}(\mathcal{A})} = \overline{\lim}_{n \rightarrow \infty} \frac{n \ln n}{-\ln \theta_{\mathcal{A}}(p, q_p, n)} \leq \rho_p(F).$$

Since $\beta_p(\mathcal{A}) = \inf_q \{\beta_{p, q}(\mathcal{A})\}$, then

$$-\frac{1}{\beta_p(\mathcal{A})} \leq -\frac{1}{\beta_{p, q_p}(\mathcal{A})} \leq \rho_p(F).$$

Hence, $\rho_p(F) \geq -\frac{1}{\beta_p(\mathcal{A})}$, $\forall p$.

Vice-versa, since

$$-\frac{1}{\beta_{p, q}(\mathcal{A})} = \overline{\lim}_{n \rightarrow \infty} \frac{n \ln n}{-\ln \theta_{\mathcal{A}}(p, q, n)},$$

then

$$\frac{n \ln n}{-\ln \theta_{\mathcal{A}}(p, q, n)} < -\frac{1}{\beta_{p, q}(\mathcal{A})} + \frac{\varepsilon}{2}, \quad \forall p \quad \forall \varepsilon > 0 \quad \forall q \quad \forall n > n_0(p, q, \varepsilon).$$

And since $\beta_p(\mathcal{A}) = \inf_q \{\beta_{p, q}(\mathcal{A})\}$, then

$$\forall p \quad \forall \varepsilon > 0 \quad \exists q_p(\varepsilon) : -\frac{1}{\beta_{p, q_p}(\mathcal{A})} \leq -\frac{1}{\beta_p(\mathcal{A})} + \frac{\varepsilon}{2}.$$

Thus,

$$\forall p \quad \forall \varepsilon > 0 \quad \exists q_p(\varepsilon) \quad \exists n_0(p, \varepsilon) \quad \forall n > n_0 : \frac{n \ln n}{-\ln \theta_{\mathcal{A}}(p, q_p, n)} < -\frac{1}{\beta_p(\mathcal{A})} + \varepsilon,$$

therefore,

$$\forall p \quad \forall \varepsilon > 0 \quad \exists q_p(\varepsilon) \quad \exists n_0(p, \varepsilon) \quad \forall n > n_0 : \sqrt[n]{\theta_{\mathcal{A}}(p, q_p, n)} < n^{-\frac{1}{-\frac{1}{\beta_p(\mathcal{A})} + \varepsilon}}.$$

By Lemma 2 $\left(b_p = -\frac{1}{\beta_p(\mathcal{A})} + \varepsilon, a_p = \frac{1}{e^{(-\frac{1}{\beta_p(\mathcal{A})} + \varepsilon)}} \right)$

$$\forall p \quad \forall \varepsilon > 0 \quad \exists q_p(\varepsilon) \quad \exists r_0(p, \varepsilon) \quad \forall r > r_0 : M_F(p, q_p, r) \leq \exp \left\{ (a_p + \varepsilon) r^{(-\frac{1}{\beta_p(\mathcal{A})} + \varepsilon)} \right\}.$$

It means that $\rho_p(F) \leq -\frac{1}{\beta_p(\mathcal{A})}$, $\forall p$.

Thus, identity (16) is proven. Identity (18) follows immediately from (16).

Let us prove identity (17).

Suppose the function F has the p -order $0 < \rho_p(F) < \infty$ and the p -type $\sigma_p(F)$. Then

$$\forall p \quad \forall \varepsilon > 0 \quad \exists q_p(\varepsilon) \quad \exists r_0(p, \varepsilon) \quad \forall r > r_0 : M_F(p, q_p, r) < \exp \{ (\sigma_p(F) + \varepsilon) r^{\rho_p(F)} \}.$$

By Lemma 1 ($a_p = \sigma_p(F) + \varepsilon$, $b_p = \rho_p(F)$) we have

$$\sqrt[n]{\theta_{\mathcal{A}}(p, q_p, n)} < \left(\frac{(\sigma_p(F) + \varepsilon) \rho_p(F) e}{n} \right)^{\frac{1}{\rho_p(F)}}, \quad \forall n > n_0,$$

$$n^{\frac{1}{\rho_p(F)}} \sqrt[n]{\theta_{\mathcal{A}}(p, q_p, n)} < ((\sigma_p(F) + \varepsilon) \rho_p(F) e)^{\frac{1}{\rho_p(F)}}, \quad \forall n > n_0.$$

By the arbitrariness of ε

$$\begin{aligned}\alpha_{p,q_p}(\mathcal{A}) &= \overline{\lim}_{n \rightarrow \infty} n^{-\beta_p(\mathcal{A})} \sqrt[n]{\theta_{\mathcal{A}}(p, q_p, n)} = \\ &= \overline{\lim}_{n \rightarrow \infty} n^{\frac{1}{\rho_p(F)}} \sqrt[n]{\theta_{\mathcal{A}}(p, q_p, n)} \leq (\sigma_p(F) \rho_p(F) e)^{\frac{1}{\rho_p(F)}}\end{aligned}$$

Since $\alpha_p(\mathcal{A}) = \inf_q \{\alpha_{p,q}(\mathcal{A})\}$, then

$$\alpha_p(\mathcal{A}) \leq \alpha_{p,q_p}(\mathcal{A}) \leq (\sigma_p(F) \rho_p(F) e)^{\frac{1}{\rho_p(F)}}, \quad \forall p.$$

Vice-versa, since

$$\alpha_{p,q}(\mathcal{A}) = \overline{\lim}_{n \rightarrow \infty} n^{-\beta_p(\mathcal{A})} \sqrt[n]{\theta_{\mathcal{A}}(p, q, n)} = \overline{\lim}_{n \rightarrow \infty} n^{\frac{1}{\rho_p(F)}} \sqrt[n]{\theta_{\mathcal{A}}(p, q, n)}, \quad \forall p, \quad \forall q,$$

then

$$\forall \varepsilon > 0 \quad \forall p \quad \exists q(p, \varepsilon) \quad \exists n_0(p, \varepsilon) \quad \forall n > n_0,$$

$$\sqrt[n]{\theta_{\mathcal{A}}(p, q, n)} < \left(\frac{(\alpha_{p,q}(\mathcal{A}) + \varepsilon)^{\rho_p(F)}}{n} \right)^{\frac{1}{\rho_p(F)}} < \left(\frac{(\alpha_p(\mathcal{A}) + 2\varepsilon)^{\rho_p(F)}}{n} \right)^{\frac{1}{\rho_p(F)}}.$$

By Lemma 2 $\left(b_p = \rho_p(F), \quad a_p = \frac{(\alpha_p(\mathcal{A}) + 2\varepsilon)^{\rho_p(F)}}{\rho_p(F)e} \right)$ we obtain

$$\forall p \quad \forall \varepsilon > 0 \quad \exists q_p(\varepsilon) \quad \exists r_0(p, \varepsilon) \quad \forall r > r_0 : \quad M_F(p, q_p, r) < \exp \{ (a_p + \varepsilon) r^{\rho_p(F)} \}.$$

It implies

$$\sigma_p(F) \leq a_p = \frac{(\alpha_p(\mathcal{A}) + 2\varepsilon)^{\rho_p(F)}}{\rho_p(F)e}.$$

By the arbitrariness of ε

$$\sigma_p(F) \rho_p(F) e \leq (\alpha_p(\mathcal{A}))^{\rho_p(F)},$$

therefore,

$$\alpha_p(\mathcal{A}) \geq (\sigma_p(F) \rho_p(F) e)^{\frac{1}{\rho_p(F)}},$$

i.e.,

$$\sigma_p(F) = -\frac{\beta_p(\mathcal{A})}{e} (\alpha_p(\mathcal{A}))^{-\frac{1}{\beta_p(\mathcal{A})}}, \quad \forall p.$$

Hence, identity (17) is proven.

Let us prove identity (19).

If $\beta_p(\mathcal{A}) < \beta(\mathcal{A})$, $\forall p$, from identity (16) it follows $\rho_p(F) < \rho(F)$, $\forall p$ and by the definition $\sigma(F) = 0$.

If $\beta_p(\mathcal{A}) = \beta(\mathcal{A})$, $\forall p$, identity (16) yields $\rho_p(F) = \rho(F)$, $\forall p$ and by the definition

$$\sigma(F) = \sup_p \{ \sigma_p(F) \} = -\frac{\beta(\mathcal{A})}{e} \sup_p \{ (\alpha_p(\mathcal{A}))^{-\frac{1}{\beta(\mathcal{A})}} \} = -\frac{\beta(\mathcal{A})}{e} (\alpha(\mathcal{A}))^{-\frac{1}{\beta(\mathcal{A})}}.$$

□

Remark. We observe that relation (16) is true also for $\rho_p(F) = \infty$. If we suppose $\rho_p(F) = \infty$ and $\beta_p(\mathcal{A}) < 0$, by (above proven) $\rho_p(F) < \infty$ that is false. Similarly, identity (17) holds also for $\sigma_p(F) = \infty$.

Remark. Formulae (16) and (17) show that p -orders and p -type of an entire operator-valued function are completely determined by the characteristics of the sequence of its coefficients.

Examples.

1. Let $\mathbf{H}_1 = \mathbf{H} = \mathbf{H}(\mathbb{C})$ be the space of all entire functions with the topology of uniform convergence on the compacts

$$\|x(z)\|_p = \max_{|z| \leq p} |x(z)|, \quad p > 0.$$

Let us find the characteristics of the function

$$F(t) = e^{t \frac{d}{dz}} = \sum_{n=0}^{\infty} \frac{t^n}{n!} \frac{d^n}{dz^n} : \mathbb{C} \rightarrow \text{Lec}(\mathbf{H}(\mathbb{C})).$$

The sequence $\mathcal{A} = \left\{ \frac{1}{n!} \frac{d^n}{dz^n} \right\}$ has the following characteristics [1],

$$\beta_p(\mathcal{A}) = \alpha_p(\mathcal{A}) = 0, \quad \forall p.$$

Therefore, $\rho_p(F) = \infty, \forall p$.

2. Let $\mathbf{H}_1 = [\rho, \sigma], \mathbf{H} = [\rho, \theta], \theta \geq \sigma$. The topologies on these spaces are determined by the multinorms

$$\|x(z)\|_\varepsilon = \sup_{p>0} \left\{ \max_{|z| \leq p} |x(z)| e^{-(\sigma+\varepsilon)p^\rho} \right\}, \quad \varepsilon > 0, \quad x \in [\rho, \sigma].$$

$$\|y(z)\|_\varepsilon = \sup_{p>0} \left\{ \max_{|z| \leq p} |y(z)| e^{-(\theta+\varepsilon)p^\rho} \right\}, \quad \varepsilon > 0, \quad y \in [\rho, \theta].$$

Let us find the characteristics of the function

$$F(t) = e^{t \frac{d}{dz}} = \sum_{n=0}^{\infty} \frac{t^n}{n!} \frac{d^n}{dz^n} : \mathbb{C} \rightarrow \text{Lec}([\rho, \sigma], [\rho, \theta]).$$

The sequence $\mathcal{A} = \left\{ \frac{1}{n!} \frac{d^n}{dz^n} \right\}$ has the following characteristics [1]:

$$\beta_\varepsilon(\mathcal{A}) = -\frac{1}{\rho}, \quad \alpha_\varepsilon(\mathcal{A}) = (\rho e \sigma \Omega_\varepsilon)^{\frac{1}{\rho}}, \quad \forall \varepsilon,$$

where

$$\Omega_\varepsilon = \begin{cases} \left(1 - \left(\frac{\sigma}{\theta+\varepsilon} \right)^{\frac{1}{\rho-1}} \right)^{1-\rho}, & \rho > 1 \\ 1, & \rho \leq 1 \end{cases}$$

Therefore,

$$\rho_\varepsilon(F) = \rho, \quad \sigma_\varepsilon(F) = \sigma \Omega_\varepsilon, \quad \forall \varepsilon.$$

3. Let $\mathbf{H}_1 = \mathbf{H} = \mathbf{H}(\mathbb{C})$ be the space of all entire functions with the topology of uniform convergence on the compacts

$$\|x(z)\|_p = \max_{|z| \leq p} |x(z)|, \quad p > 0.$$

Let us find the characteristics of the function

$$\begin{aligned} F(t)(x) &= x(z) + t \int_0^z e^{(z-\xi)t} x(\xi) d\xi = \\ &= x(z) + t \sum_{n=0}^{\infty} \int_0^z \frac{(z-\xi)^n t^n}{n!} x(\xi) d\xi, \end{aligned}$$

$$F(t) : \mathbb{C} \rightarrow \text{Lec}(\mathbf{H}(\mathbb{C})).$$

Here

$$A_n(x) = \int_0^z \frac{(z-\xi)^{n-1}}{(n-1)!} x(\xi) d\xi, \quad n = 1, 2, \dots, \quad A_0 = E.$$

The sequence $\mathcal{A} = \{A_n\}$ has the following characteristics [1],

$$\beta_p(\mathcal{A}) = -1, \quad \alpha_p(\mathcal{A}) = p, \quad \forall p.$$

Therefore, $\rho_p(F) = 1, \quad \sigma_p(F) = \frac{p}{e}, \quad \forall p.$

3. PROPERTIES OF GROWTH CHARACTERISTICS FOR OPERATOR-VALUED FUNCTIONS

Let us note certain properties of the growth characteristics for operator-valued functions implied by Theorem 4.

1⁰. Entire function F and its k th derivative $F^{(k)}$ has the same p -orders and p -types of growth.

The validity follows from the fact the sequences $\{A_n\}$ and $\left\{\frac{(n+k)!}{n!}A_{n+k}\right\}$ has the same characteristics for each fixed k .

2⁰. If a function F_1 has the p -orders $\rho_p(F_1)$ and the p -types $\sigma_p(F_1)$, and a function F_2 has the p -orders $\rho_p(F_2) > \rho_p(F_1), \forall p$ and the p -types $\sigma_p(F_2)$, the function $F = F_1 + F_2$ has the p -orders $\rho_p(F) = \rho_p(F_2), \forall p$ and the p -types $\sigma_p(F) = \sigma_p(F_2), \forall p$.

The validity is implied by the fact that the characteristics of the sum of operators are equal to the characteristics of the term of the greater order.

3⁰. If a function F_1 has the p -orders $\rho_p(F_1)$ and the p -types $\sigma_p(F_1)$, and a function F_2 has the p -orders $\rho_p(F_2) = \rho_p(F_1), \forall p$ and the p -types $\sigma_p(F_2) > \sigma_p(F_1), \forall p$, then the function $F = F_1 + F_2$ has the p -orders $\rho_p(F) = \rho_p(F_2), \forall p$ and the p -types $\sigma_p(F) = \sigma_p(F_2), \forall p$.

The validity follows from the fact that the characteristics of the sum of operators of same orders are equal to the characteristics of the term of the greater type.

4⁰. (The case $\mathbf{H}_1 = \mathbf{H}$.) Suppose a function F_1 has the order $\rho(F_1)$ and the type $\sigma(F_1)$, and a function F_2 has the order $\rho(F_2) > \rho(F_1)$ and the type $\sigma(F_2)$. Then the function $F = F_1F_2$ has the order $\rho(F) \leq \rho(F_2)$ and the type $\sigma(F)$. If $\rho(F) = \rho(F_2)$, then $\sigma(F) \leq \sigma(F_2)$. A similar statement holds for the function $\tilde{F} = F_2F_1$.

The proof is based on the following lemma.

Lemma 3. Suppose a sequence of operators $\mathcal{A} = \{A_n\}$ has the order $\beta(\mathcal{A})$ and the type $\alpha(\mathcal{A})$, and a sequence of the operators $\mathcal{B} = \{B_n\}$ has the order $\beta(\mathcal{B}) > \beta(\mathcal{A})$ and the type $\alpha(\mathcal{B})$. Then the sequence of the operators $\mathcal{C} = \{C_n\}$, where $C_n = \sum_{k=0}^n A_k B_{n-k}$ has the order $\beta(\mathcal{C}) \leq \beta(\mathcal{B})$ and the type $\alpha(\mathcal{C})$. If $\beta(\mathcal{C}) = \beta(\mathcal{B})$, then $\alpha(\mathcal{C}) \leq \alpha(\mathcal{B})$.

Proof. Denote $a = \alpha(\mathcal{A})e^{\beta(\mathcal{A})}, \quad b = \alpha(\mathcal{B})e^{\beta(\mathcal{B})}$.

The definition of the order and type of a sequence of operators implies [1]

$$\forall \varepsilon, \varepsilon_1 > 0, \quad \forall p, \quad \exists M_p, \quad \exists q, \quad \forall n, \quad \forall x \in \mathbf{H} :$$

$$\begin{aligned} \|C_n(x)\|_p &\leq M_p \left((b + \varepsilon)^n n!^{\beta(\mathcal{B})} + (a + \varepsilon_1)(b + \varepsilon)^{n-1} 1!^{\beta(\mathcal{A})} (n-1)!^{\beta(\mathcal{B})} + \dots + \right. \\ &\quad \left. + (a + \varepsilon_1)^{n-1} (b + \varepsilon)(n-1)!^{\beta(\mathcal{A})} 1!^{\beta(\mathcal{B})} + (a + \varepsilon_1)^n n!^{\beta(\mathcal{A})} \right) \|x\|_q \leq \\ &\leq M_p (b + \varepsilon)^n n!^{\beta(\mathcal{B})} \left[1 + \binom{n}{1}^{-\beta(\mathcal{B})} \left(\frac{a + \varepsilon_1}{b + \varepsilon} \right) 1!^\nu + \binom{n}{2}^{-\beta(\mathcal{B})} \left(\frac{a + \varepsilon_1}{b + \varepsilon} \right)^2 2!^\nu + \dots + \right. \\ &\quad \left. + \binom{n}{n}^{-\beta(\mathcal{B})} \left(\frac{a + \varepsilon_1}{b + \varepsilon} \right)^n n!^\nu \right] \|x\|_q, \quad (21) \end{aligned}$$

where $\nu = \beta(\mathcal{A}) - \beta(\mathcal{B})$.

If $\beta(\mathcal{B}) > \beta(\mathcal{A})$ ($\nu < 0$), for large n the expression in brackets in (21) does not exceed $(1 + \varepsilon_2)^n$, $\forall \varepsilon_2 > 0$ and thus $\beta(\mathcal{C}) \leq \beta(\mathcal{B})$, and if $\beta(\mathcal{C}) = \beta(\mathcal{B})$, then $\alpha(\mathcal{C}) \leq \alpha(\mathcal{B})$. \square

5⁰. (The case $\mathbf{H}_1 = \mathbf{H}$.) Suppose a function F_1 has the order $\rho(F_1)$ and the type $\sigma(F_1)$, and a function F_2 has the order $\rho(F_2) = \rho(F_1)$ and the type $\sigma(F_2) \geq \sigma(F_1)$. Then the function $F = F_1 F_2$ has the order $\rho(F) \leq \rho(F_2)$ and the type $\sigma(F)$. If $\rho(F) = \rho(F_2)$, then $\sigma(F) \leq 2\sigma(F_2)$. A similar statement holds true for the function $\tilde{F} = F_2 F_1$.

The proof is based on the following lemma.

Lemma 4. Let a sequence of operators $\mathcal{A} = \{A_n\}$ has the order $\beta(\mathcal{A})$ and the type $\alpha(\mathcal{A})$, and a sequence of operators $\mathcal{B} = \{B_n\}$ does the order $\beta(\mathcal{B}) = \beta(\mathcal{A})$ and the type $\alpha(\mathcal{B}) \geq \alpha(\mathcal{A})$. The the sequence of the operators $\mathcal{C} = \{C_n\}$, where $C_n = \sum_{k=0}^n A_k B_{n-k}$, has the order $\beta(\mathcal{C}) \leq \beta(\mathcal{B})$ and the type $\alpha(\mathcal{C})$. If $\beta(\mathcal{C}) = \beta(\mathcal{B})$, then $\alpha(\mathcal{C}) \leq 2^{-\beta(\mathcal{B})} \alpha(\mathcal{B})$.

Proof. Under the hypothesis of the lemma the expression in the brackets in (21) does not exceed $2^{-\beta(\mathcal{B})n} n$ and thus $\beta(\mathcal{C}) \leq \beta(\mathcal{B})$, and if $\beta(\mathcal{C}) = \beta(\mathcal{B})$, then $\alpha(\mathcal{C}) \leq 2^{-\beta(\mathcal{B})} \alpha(\mathcal{B})$. \square

Remark. As it is known, in the scalar case the theorem on categories [3, Th. 12] is valid. In the case of operator-valued function this question is still open.

6⁰. (Invariance). Suppose $\mathbf{H}_1, \tilde{\mathbf{H}}_1, \mathbf{H}$ and $\tilde{\mathbf{H}}$ are four locally convex spaces with the topologies induced respectively by the multinorms $\|\cdot\|'_q, q \in \mathcal{Q}, \|\cdot\|'_q, \tilde{q} \in \tilde{\mathcal{Q}}, \|\cdot\|_p, p \in \mathcal{P}, \|\cdot\|_{\tilde{p}}, \tilde{p} \in \tilde{\mathcal{P}}$ and let $T_1 : \mathbf{H}_1 \rightarrow \tilde{\mathbf{H}}_1, T : \mathbf{H} \rightarrow \tilde{\mathbf{H}}$ are two topological isomorphisms. Then

1) for each operator-valued function

$$F(t) = \sum_{n=0}^{\infty} A_n t^n : \mathbb{C} \rightarrow \text{Lec}(\mathbf{H}_1, \mathbf{H})$$

its order and type coincide with the order and type of the function

$$\tilde{F}(t) = \sum_{n=0}^{\infty} T A_n T_1^{-1} t^n : \mathbb{C} \rightarrow \text{Lec}(\tilde{\mathbf{H}}_1, \tilde{\mathbf{H}});$$

2) if all the p -orders of the function F are strictly less than its order, then all the \tilde{p} -orders of the function \tilde{F} are strictly less than its order;

3) if at least one p -order of the function F equals to its order, then at least one \tilde{p} -order of the function \tilde{F} equals to its order;

4) if the function F has the p -orders $\rho_p(F)$, the order $\rho(F)$, the p -types $\sigma_p(F)$ and the type $\sigma(F)$, at that the set

$$\mathcal{P}_F = \{p \in \mathcal{P} : \rho_p(F) = \rho(F)\}$$

is non-empty and $\forall p \in \mathcal{P}_F : \sigma_p(F) < \sigma(F)$, then the function \tilde{F} has the \tilde{p} -orders $\rho_{\tilde{p}}(\tilde{F})$, the order $\rho(\tilde{F})$, the \tilde{p} -types $\sigma_{\tilde{p}}(\tilde{F})$ and the type $\sigma(\tilde{F})$, at that the set

$$\tilde{\mathcal{P}}_{\tilde{F}} = \{\tilde{p} \in \tilde{\mathcal{P}} : \rho_{\tilde{p}}(\tilde{F}) = \rho(\tilde{F})\}$$

is non-empty and $\forall \tilde{p} \in \tilde{\mathcal{P}}_{\tilde{F}} : \sigma_{\tilde{p}}(\tilde{F}) < \sigma(\tilde{F})$;

5) if under hypothesis of Item 4) $\exists p \in \mathcal{P}_F : \sigma_p(F) = \sigma(F)$, then $\exists \tilde{p} \in \tilde{\mathcal{P}}_{\tilde{F}} : \sigma_{\tilde{p}}(\tilde{F}) = \sigma(\tilde{F})$.

The validity of property 6⁰ follows from analogous properties for the characteristics of a sequence of operators [1, 6].

The invariance property implies that under any replacements of the multinorms in \mathbf{H}_1 and \mathbf{H} to equivalent ones (T_1 and T are identity operators)

1) the order and type of the operator-valued function F remain the same;

2) if all the p -orders of the function F were strictly less than its order before the replacement of the multinorms, after the replacement of the multinorms all its \tilde{p} -orders are also strictly less than the order;

3) if at least one p -order of the function F equals its order before the replacement of the multinorms, after the replacement at least one its \tilde{p} -order (not necessarily the same) is also equal to its order;

4) if the function F has the p -order $\rho_p(F)$ before the replacement of the multinorms, the order $\rho(F)$, the p -types $\sigma_p(F)$, and the type $\sigma(F)$, at that the set

$$\mathcal{P}_F = \{p \in \mathcal{P} : \rho_p(F) = \rho(F)\}$$

is non-empty and $\forall p \in \mathcal{P}_F : \sigma_p(F) < \sigma(F)$, then after the replacement of the multinorms this function has the \tilde{p} -orders $\rho_{\tilde{p}}(F)$, the order $\rho(F)$, the \tilde{p} -types $\sigma_{\tilde{p}}(F)$, and the type $\sigma(F)$, at that the set

$$\tilde{\mathcal{P}}_F = \{\tilde{p} \in \tilde{\mathcal{P}} : \rho_{\tilde{p}}(F) = \rho(F)\}$$

is non-empty and $\forall \tilde{p} \in \tilde{\mathcal{P}}_F : \sigma_{\tilde{p}}(F) < \sigma(F)$;

5) if under the hypothesis of Item 4) $\exists p \in \mathcal{P}_F : \sigma_p(F) = \sigma(F)$, then $\exists \tilde{p} \in \tilde{\mathcal{P}}_F : \sigma_{\tilde{p}}(F) = \sigma(F)$.

BIBLIOGRAPHY

1. V.P. Gromov, S.N. Mishin, S.V. Panyushkin. *Operators of finite order and differential-operator equations*. Orel State University, Orel. 2009. (in Russian).
2. L.V. Kantorovich, G.P. Akilov. *Functional analysis in normed spaces*. Fizmatgiz, Moscow. 1959. [International Series of Monographs in Pure and Applied Mathematics, V. 46. Pergamon Press, N.Y. 1964.]
3. B.Ya. Levin. *Distribution of Zeros of Entire Functions*. GITTL, Moscow. 1956. [American Mathematical Soc., Translations of mathematics monographs. V. 5. 1980.]
4. A.F. Leont'ev. *Entire functions. Exponential series*. Nauka, Moscow. 1983. (in Russian).
5. S.N. Mishin. *Connection of the characteristics of sequence of the operators with convergence by bornology*. // Vestnik RUDN. Ser. matem. informat. fiz. 2010. No. 4. P. 26-34. (in Russian).
6. S.N. Mishin. *Invariance of characteristics of sequence of operators and operator characteristics of vector*. // Uchenye zapiski OGU (lab. TFFA). 2010. No. 7. P. 32-35. (in Russian).
7. S.N. Mishin. Finite order operators in locally convex spaces and there applications. Dissertation for candidate of phys. and math. sci. degree. Orel. 2002.
8. Ya.V. Radyno. *Linear equations and bornology*. Belorussian State University, Minsk. 1982.
9. P. Garrett. *Holomorphic vector-valued functions*. Course notes. 2005. Electronic resource. URL: http://www.math.umn.edu/~garrett/m/fun/Notes/09_vv_holo.pdf

Sergei Nikolaevich Mishin,
 Orel State University,
 Komsomolskaya str., 95,
 302026, Orel, Russia
 E-mail: sergeymishin@rambler.ru