

ANALYTIC FUNCTIONS WITH SMOOTH ABSOLUTE VALUE OF BOUNDARY DATA

F.A. SHAMOYAN

Abstract. Let f be an analytic function in the unit circle D continuous up to its boundary Γ , $f(z) \neq 0$, $z \in D$. Assume that on Γ , the function $|f|$ has a modulus of continuity $\omega(|f|, \delta)$. In the paper we establish the estimate $\omega(f, \delta) \leq A\omega(|f|, \sqrt{\delta})$, where A is a some non-negative number, and we prove that this estimate is sharp. Moreover, in the paper we establish a multi-dimensional analogue of the mentioned result. In the proof of the main theorem, an essential role is played by a theorem of Hardy-Littlewood type on Hölder classes of the functions analytic in the unit circle.

Keywords: analytic function, modulus of continuity, factorization, outer function.

Mathematics Subject Classification: primary: 30D55, 30D15; secondary: 46E22, 47A15

INTRODUCTION

Let $D = \{z \in \mathbb{C} : |z| < 1\}$ be a bounded unit circle in the complex plane \mathbb{C} and Γ be its boundary. By C_A we denote the set of all functions f analytic in D and continuous in $D \cup \Gamma$. If $f \in C(\Gamma)$, then by the symbol $\omega(f, \delta)$ we denote the modulus of continuity of the function f on Γ , that is,

$$\omega(f, \delta) = \left\{ \sup_{\gamma \in \Gamma, |t| \leq \delta, t \in \mathbb{R}} |f(\gamma) - f(\gamma e^{it})| \right\}.$$

In the paper we consider the following problem: let $f \in C_A$ and at the unit circle the function $|f(e^{i\theta})|$ has the modulus of continuity $\omega(|f|, \delta)$. What is the modulus of continuity of the function f on Γ and hence, on $D \cup \Gamma$?

Such problem in the classes of continuous functions with the modulus of continuity satisfying Bari-Stechkin condition

$$\int_0^\delta \frac{\omega(|f|, t)}{t} dt + \delta \int_\delta^\pi \frac{\omega(|f|, t)}{t^2} dt = O(\omega(|f|, \delta)), \quad \delta \rightarrow 0, \quad (1)$$

was solved first in work by V.P. Khaving and the author, see [5].

It was established that if $\omega(|f|, \delta)$ satisfies Bari-Stechkin condition (1) and $f(z) \neq 0$, $z \in D$, then

$$\omega(f, \delta) = O(\omega(|f|, \sqrt{\delta})), \quad \delta \rightarrow 0.$$

Moreover, it was shown by simple examples that the obtained estimate was sharp and the condition $f(z) \neq 0$, $z \in D$, is necessary in the known sense. The detailed proof of these statements was exposed in [7]. This work gave rise to rather interesting studies in this direction. First V.P. Khavin, see [6], proposed an interesting approach for obtaining such estimates; this was done by applying the methods of the theory of singular integral operators. Later N.A. Shirokov (see [8], [10]) extended the results of such type for external functions and Hölder

F.A. SHAMOYAN, ANALYTIC FUNCTIONS WITH SMOOTH ABSOLUTE VALUE OF BOUNDARY DATA.

© SHAMOYAN F.A. 2017.

Submitted May 10, 2017.

classes of order $\alpha, \alpha \in (0, +\infty)$ and he obtained the necessary and sufficient condition for $|f(e^{i\theta})|$ ensuring that the function f has a prescribed modulus of continuity on the set $D \cup \Gamma$. In these works there were introduced a new characteristics and in terms of this characteristics, N.A. Shirokov obtained the results of such kind also for the Besov classes of analytic functions in $D \cup \Gamma$. And finally, we mention work [2], where it was established that this phenomenon has a local character, that is, if the modulus of continuity $|f|$ on the circle satisfies the Hölder condition of order α just at one point, then f belongs to the Hölder class of order $\frac{\alpha}{2}$ at this point.

We note that the proof by V.P. Khavni and the proof of the results in works [2], [6], [8], [10] are based on gentle and fine methods of complex and harmonic analysis. In our opinion, the approach applied in works [5] and [7] and is based on classical theorems of Hardy-Littlewood type theorems (see [3], [4]) is more simple. In this work by developing the methods of works [5], [7] we prove such results for the modulus of continuity $\omega(|f|, \delta)$ satisfying the classical Zygmund condition

$$\int_0^\delta \frac{\omega(|f|, t)}{t} dt = O(\omega(|f|, \delta)), \quad \delta \rightarrow 0. \quad (2)$$

The following statement is true.

Theorem 1. *Let f be a function in the class C_A and $f(z) \neq 0, z \in D$. If the modulus of the continuity $\omega(|f|, \delta)$ of the function $|f|$ on Γ satisfies Zygmund condition (2), then*

$$\omega(f, \delta) = O\left(\omega(|f|, \sqrt{\delta})\right), \quad \delta \rightarrow 0, \quad (3)$$

and this estimate is sharp.

Remark 1. *A simple example, the function*

$$f(z) = (1 - z)^{2\alpha} \exp\left(-\frac{1 + z}{1 - z}\right), \quad z \in D \cup \Gamma, \quad \alpha \in (0, +\infty),$$

where the principal branch of the power function is chosen, shows the sharpness of the statement of Theorem 1.

The analogue of Theorem 1 is true for analytic functions in the unit ball of the space \mathbb{C}^n . In order to formulate it, we introduce some notations. Let $z = (z_1, \dots, z_n) \in \mathbb{C}^n, \|z\| = \sqrt{|z_1|^2 + \dots + |z_n|^2}$. We define $B_n = \{z \in \mathbb{C}^n : \|z\| < 1\}$ and $S_n = \{z \in \mathbb{C}^n : \|z\| = 1\}$.

By $H(B_n)$ we denote the set of all analytic functions in B_n . Let $f \in H(B_n)$ and $f(z) = \sum_{k=0}^{+\infty} f_k(z)$ be the expansions of the function f into homogeneous polynomials; by $R(f)$ the radial derivative of the function f [11], that is,

$$R(f)(z) = \sum_{k=1}^{+\infty} k f_k(z), \quad z \in B_n.$$

We introduce also the notation:

$$C_A(B_n) = H(B_n) \cap C(B_n \cup S_n).$$

The following estimate of Theorem 1 is true in the classes $C_A(B_n)$.

Theorem 2. *Let $f \in C_A(B_n)$ and the modulus of continuity $\omega(|f|, \delta)$ of the function $|f|$ on S_n satisfies Zygmund condition (2). Then the modulus of continuity of the function on the set $B_n \cup S_n$ satisfies the estimate*

$$\omega(f, \delta) \leq A\omega(|f|, \sqrt{\delta}), \quad 0 \leq \delta \leq 2.$$

Remark 2. For Hölder classes, that is, as $\omega(f, t) = t^\alpha, 0 < \alpha \leq 1, t \in [0, 2]$, the analogue of Theorem 2 was established in work [9] by N.A. Shirokov.

1. PROOF OF AUXILIARY STATEMENTS

Let f and g be real-valued functions with a common domain $E \subset \mathbb{C}$. Then the relation $f \lesssim g$ on E is equivalent to the following: there exists a positive number A such that $f(z) \leq Ag(z)$ for each $z \in E$. If $f \lesssim g$ and simultaneously $g \lesssim f$, then $f(z) \approx g(z)$.

In what follows, as a function of modulus of continuity type we call a non-negative non-decreasing function ω on $[0, +\infty)$ such that

$$\omega(0) = 0, \quad \omega(\delta_1 + \delta_2) \leq \omega(\delta_1) + \omega(\delta_2), \quad \omega(\lambda\delta) \leq 2\lambda\omega(\delta), \quad \lambda, \delta \in [0, +\infty).$$

Lemma 1. Let ω be a function of modulus of continuity type satisfying Zygmund condition (2), then

$$\omega(\delta) \ln \frac{1}{\delta} \lesssim \omega(\sqrt{\delta}), \quad \delta > 0. \quad (4)$$

Proof. By the definition we have

$$\int_0^{\sqrt{\delta}} \frac{\omega(u)}{u} du \leq A\omega(\sqrt{\delta}).$$

It is clear that if $1 \leq \delta$, then estimate (4) is obvious and this is why we assume that $0 < \delta < 1$. Then

$$\int_0^{\sqrt{\delta}} \frac{\omega(u)}{u} du = \int_0^{\delta} \frac{\omega(u)}{u} du + \int_{\delta}^{\sqrt{\delta}} \frac{\omega(u)}{u} du.$$

Hence,

$$\int_0^{\sqrt{\delta}} \frac{\omega(u)}{u} du \geq \omega(\delta) \int_{\delta}^{\sqrt{\delta}} \frac{du}{u} = \frac{\omega(\delta)}{2} \ln \frac{1}{\delta}.$$

It remains to employ Zygmund condition. The proof is complete. \square

Lemma 2. Let $f \in C_A, t = |t|\tau, t \in D, \tau \in \Gamma$. Then the estimate

$$|f(t)| \lesssim \left(|f(\tau)| + \omega(f, (1 - |t|)) \ln \frac{1}{1 - |t|} \right)$$

is true.

Proof. We have

$$f(t) = \frac{1}{2\pi} \int_{\Gamma} P_t(\xi) f(\xi) |d\xi|.$$

where $P_t(\xi)$ is the Poisson kernel. This is why

$$|f(t)| \leq \frac{1}{2\pi} \int_{\Gamma} P_t(\xi) |f(\xi) - f(\tau)| |d\xi| + |f(\tau)|.$$

Therefore,

$$|f(t)| \lesssim (|f(\tau)| + J_\omega), \quad (5)$$

where

$$J_\omega := \frac{1}{2\pi} \int_{\Gamma} P_t(\xi) \omega(f, |\xi - \tau|) d\tau.$$

We proceed to estimating the latter integral. It is clear that

$$J_\omega \lesssim \int_{\Gamma} \frac{(1-|t|^2)\omega(f, |\xi-\tau|)}{(1-|t|)^2 + |\xi-\tau|^2} |d\xi| \lesssim \int_0^\pi \frac{(1-|t|^2)\omega(f, u)}{(1-|t|)^2 + u^2} du.$$

We define $\omega(f, \delta)$ on $\mathbb{R}_+ = [0, +\infty)$ as a function of modulus of continuity type (see [1], [4]). Then

$$J_\omega \lesssim \int_0^{\frac{\pi}{1-|t|}} \frac{\omega(f, v(1-|t|))}{1+v^2} dv.$$

Representing this integral as the sum

$$\int_0^1 \frac{\omega(f, v(1-|t|))}{1+v^2} dv + \int_1^{\frac{\pi}{1-|t|}} \frac{\omega(f, v(1-|t|))}{1+v^2} dv,$$

and taking into consideration that $\frac{\omega(f, \delta)}{\delta}$ does not increase (see [1], [4]), we obtain

$$J_\omega \lesssim \omega(f, 1-|t|) \int_1^{\frac{\pi}{1-|t|}} \frac{v}{1+v^2} dv.$$

Hence,

$$J_\omega \lesssim \omega(f, 1-|t|) \ln \frac{1}{1-|t|}. \quad (6)$$

By (5), (6) we arrive at the statement of the lemma. \square

Lemma 3. (see [3]). Let λ be a positive non-decreasing function on $(0, 1)$ and

$$\int_0^1 \lambda(u) du < +\infty.$$

Assume that f is an analytic function in D such that

$$\sup_{z \in D} \left\{ \frac{|f'(z)|}{\lambda(|z|)} \right\} < +\infty.$$

Then the function f belongs to the class C_A and

$$\omega(f, \delta) \lesssim \int_{1-\delta}^1 \lambda(u) du.$$

Lemma 4. Let $f \in C_A$, $f(z) \neq 0$, $z \in D$, $|f(z)| \leq 1$, $z \in D$. Then there exists a number $M > 0$ possessing the following properties: for an arbitrary $0 < a < 1$, the function f in D can be represented as

$$f(z) = \Phi_a(z)\Psi_a(z), \quad z \in D,$$

where Ψ_a is an analytic function in D such that $|\Psi_a|$ is continuously extended to the entire closed circle $D \cup \Gamma$,

$$\begin{aligned} a &\leq |\Psi_a(t)| \leq 1 \quad \text{for each } t \in D, \\ ||\Psi_a(t')| - |\Psi_a(t'')|| &\leq ||f(t')| - |f(t'')||, \quad t', t'' \in \Gamma, \\ \int_{\Gamma} |\ln |\Psi_a(t)|| |dt| &\leq M, \\ \Phi_a(z) &= \exp \left(- \int_{\Gamma} S_z(t) d\mu_a(t) \right), \quad z \in D, \end{aligned} \tag{7}$$

where $S_z(t)$ is the Schwarz kernel for the circle D , μ_a is a non-negative Borel measure Γ , whose total variation does not exceed M .

Proof. Let $H_a(t) = \max(a, |f(t)|)$, $t \in \Gamma$. We have

$$\begin{aligned} \Psi_a(z) &:= \exp \left(\frac{1}{2\pi} \int_{\Gamma} S_z(t) \ln H_a(t) |dt| \right), \\ \Phi_a(z) &= f(z) (\Psi_a(z))^{-1} := \exp \left(- \int_{\Gamma} S_z(t) d\mu_a(t) \right), \quad z \in D, \end{aligned}$$

where

$$\mu_a(E) = - \int_{E_a} \ln \left(\frac{|f(t)|}{a} \right) |dt| + \mu(E), \tag{8}$$

μ is a non-negative measure defining the singular part of the function f , E is an arbitrary Borel set in Γ , $E_a = \{\gamma \in \Gamma : |f(\gamma)| \leq a\}$. It is clear that $|\Psi_a|$ on Γ coincides with H_a . Estimate (7) is obtained by the following inequality

$$\begin{aligned} \int_{\Gamma} |\ln |\Psi_a(\xi)|| |d\xi| &= \int_{E(|f| \geq a)} |\ln |\Psi_a(\xi)|| |d\xi| + \int_{E(|f| < a)} |\ln |\Psi_a(\xi)|| |d\xi| \\ &= \int_{E(|f| \geq a)} |\ln |\Psi_a(\xi)|| |d\xi| + \sigma(E_a) \ln \frac{1}{a}, \end{aligned}$$

where σ is the Lebesgue measure on E_a .

It remains to note that

$$E_a = E(|f| \leq a) = E \left(\ln \frac{1}{|f|} \geq \ln \frac{1}{a} \right);$$

re recall that $\max |f| \leq 1$, $0 < a < 1$.

The finiteness of the integral $\int_{\Gamma} |\ln |f(\xi)|| |d\xi|$ implies that

$$\sup_{A \geq 0} A \sigma(\gamma \in \Gamma : |\ln |f(\gamma)|| \geq A) < +\infty.$$

This proves (7).

Now we are going to estimate $\mu_a(E)$. In order to do it, we denote by $V_a(E)$, the first term in the right hand side in (8) and we note that $\mu_a(\Gamma) \leq V_a(\Gamma) + \mu(\Gamma)$. This is why

$$V_a(\Gamma) \leq \int_{\Gamma} |\ln |f|| |dt| + |\ln a| \sigma(\Gamma_a),$$

where

$$\Gamma_a = \{\gamma \in \Gamma : |\ln |f(\gamma)|| \geq |\ln a|\}.$$

Following the lines of the proof of inequality (7), we obtain the last statement of the lemma. The proof is complete. \square

Remark 3. *Employing the Jensen inequality, it is easy to observe that if $\|f\|_{C_A} < 1$, then*

$$|\mu_a(\Gamma)| \lesssim \ln \frac{1}{|f(0)|}, \quad \int_{\Gamma} |\ln |\Psi_a|| |dt| \lesssim \ln \frac{1}{|f(0)|}.$$

2. PROOF OF THEOREMS

Proof of Theorem 1. Without loss of generality we assume that $|f(t)| \leq 1$, $t \in \Gamma$. Moreover, for the sake of convenience we denote $\omega(\delta) := \omega(|f|, \delta)$, $0 \leq \delta \leq 2$, and at that,

$$|f(t')| - |f(t'')| \leq \frac{1}{2}\omega(|t' - t''|) \quad \text{for all } t', t'' \in \Gamma.$$

Employing Lemma 3, it is sufficient to establish the estimate

$$|f'(t)| \lesssim \frac{\omega(\sqrt{1-|t|})}{1-|t|}, \quad t \in D.$$

Let $t \in D$ be a fixed point in the circle D and in Lemma 4 we choose $a = \omega(\sqrt{1-|t|})$. We introduce the notations

$$F_t(t) = \Psi_{\omega(\sqrt{1-|t|})}(t), \quad f_t(t) = \Phi_{\omega(\sqrt{1-|t|})}(t).$$

We observe that

$$f'(t) = f'_t(t)F'_t(t) + f'_t(t)F_t(t)$$

1°. *Estimate for $|f_t(t)||F'_t(t)|$.* To estimate this product, let us first prove the inequality

$$|F_t(t)| \lesssim |F_t(\tau)|, \quad t = |t|\tau, \quad \tau \in \Gamma. \quad (9)$$

By Lemma 2,

$$|F_t(t)| \lesssim \left(|F_t(\tau)| + \omega(1-|t|) \ln \frac{1}{1-|t|} \right).$$

This is why to prove inequality (9), it is sufficient to establish the estimate

$$\sup_{t \in D} \left\{ \omega(1-|t|) \ln \frac{1}{1-|t|} |F_t^{-1}(\tau)| \right\} < +\infty.$$

Suppose first that

$$\max(|f(\tau)|, \omega(\sqrt{1-|t|})) = |f(\tau)|,$$

then by Lemma 4 we have

$$|F_t(\tau)| \geq \omega(\sqrt{1-|t|}).$$

This is why, taking into consideration the estimate

$$\omega(1-|t|) \ln \frac{1}{1-|t|} \lesssim \omega(\sqrt{1-|t|}),$$

we obtain

$$\frac{\omega(1-|t|) \ln \frac{1}{1-|t|}}{|F_t(\tau)|} \lesssim \frac{\omega(\sqrt{1-|t|})}{\omega(\sqrt{1-|t|})} \lesssim 1.$$

Now we consider the case $|f(\tau)| \leq \omega(\sqrt{1-|t|})$. Applying Lemma 1 once again, we get desired estimate (9).

We proceed to estimating the functions $|f_t(t)|$, $|F'_t(t)|$. Let

$$\Gamma_1 = \{\gamma \in \Gamma : \omega(|\gamma - \tau|) \leq |F_t(\tau)|\}, \quad \Gamma_2 = \Gamma \setminus \Gamma_1.$$

Then we have

$$|f_t(t)| |F'_t(t)| \leq |F'_t(t)|.$$

At that,

$$\begin{aligned} |F'_t(t)| &= |F_t(t)| \left| \int_{\Gamma} |F_t(\gamma)| \frac{2\gamma}{(\gamma - t)^2} |d\gamma| \right| = |F_t(t)| \left| \int_{\Gamma} \frac{(\ln |F_t(\gamma)| - \ln |F_t(\tau)|)}{(\gamma - t)^2} \right| \\ &\lesssim |F_t(\tau)| \left[\left| \int_{\Gamma_1} \frac{(\ln |F_t(\gamma)| - \ln |F_t(\tau)|)}{(\gamma - t)^2} \right| + \left| \int_{\Gamma_2} \frac{(\ln |F_t(\gamma)| - \ln |F_t(\tau)|)}{(\gamma - t)^2} \right| \right] \\ &\lesssim |F_t(\tau)| \int_{\Gamma_1} \frac{|\ln |F_t(\gamma)| - \ln |F_t(\tau)||}{|\gamma - t|^2} |d\gamma| + |F_t(\tau)| \int_{\Gamma_2} \frac{|\ln |F_t(\gamma)||}{|\gamma - t|^2} |d\gamma| \\ &\quad + |F_t(\tau)| |\ln |F_t(\tau)|| \int_{\Gamma_2} \frac{|d\gamma|}{|\gamma - t|^2} \stackrel{def}{=} [I_1 + I_2 + I_3]. \end{aligned}$$

Estimate for I_1 . If $\gamma \in \Gamma_1$, by the mean theorem we have

$$|\ln |F_t(\gamma)|| - |\ln |F_t(\tau)|| \leq \frac{||F_t(\gamma)| - |F_t(\tau)||}{\min_{\gamma \in \Gamma_1} (|F_t(\gamma)|, |F_t(\tau)|)} \leq \frac{||f(\gamma)| - |f(\tau)||}{\min_{\gamma \in \Gamma_1} (|F_t(\gamma)|, |F_t(\tau)|)}.$$

In view of the definition of Γ_1 , we have

$$|F_t(\gamma)| \geq ||F_t(\tau)| - |F_t(\tau) - F_t(\gamma)|| \geq |F_t(\tau)| - \frac{1}{2}\omega(|\gamma - \tau|) \geq \frac{1}{2}|F_t(\tau)|.$$

Therefore,

$$I_1 \lesssim \frac{1}{1 - |t|} \int_{\Gamma} \omega(|\gamma - \tau|) P_t(\gamma) |d\gamma|,$$

where $P_t(\gamma)$ is the Poisson kernel.

Employing Lemma 2, we obtain

$$I_1 \lesssim \frac{\omega(1 - |t|)}{1 - |t|} \ln \frac{1}{1 - |t|}.$$

By Lemma 1, we finally get

$$I_1 \lesssim \frac{\omega(\sqrt{1 - |t|})}{1 - |t|}, \quad t \in D.$$

Estimate for I_2 . Let

$$K_t(\gamma) = \frac{1}{(\gamma - t)^2}, \quad \gamma \in \Gamma, \quad t \in D.$$

Then we have

$$I_2 \leq |F_t(\tau)| \max_{\gamma \in \Gamma_2} |K_t(\gamma)| \int_{\Gamma} |\ln |f_t(\gamma)|| |d\gamma| \lesssim |F_t(\tau)| \max_{t \in \Gamma_2} |K_t(\gamma)|.$$

In the latter estimate we have employed Lemma 4.

Now, taking into consideration the definition of Γ_2 , we obtain

$$I_2 \lesssim |F_t(\tau)| \max_{\gamma \in \Gamma_2} \frac{1}{(|\gamma - \tau|^2 + (1 - |t|))^2} \lesssim |F_t(\tau)| \max \left\{ \frac{1}{x^2}, x : \omega(x) \geq |F_t(\tau)| \right\}.$$

Let $x^* \in (0, 2]$ be such that

$$\omega(x^*) = |F_t(\tau)|. \tag{10}$$

Then by the latter estimate we get

$$I_2 \lesssim \frac{|F_t(\tau)|}{(x^*)^2} = C_f \frac{\omega(x^*)}{(x^*)^2}.$$

The inequality $|F_t(\tau)| \geq \omega(\sqrt{1-|t|})$ implies $\sqrt{1-|t|} \leq x^*$. This is why,

$$\frac{\omega(x^*)}{(x^*)^2} = \frac{\omega(x^*)}{x^* \cdot x^*} \leq \frac{\omega(\sqrt{1-|t|})}{\sqrt{1-|t|}} \frac{1}{\sqrt{1-|t|}} = \frac{\omega(\sqrt{1-|t|})}{1-|t|},$$

that is,

$$I_2 \lesssim \frac{\omega(\sqrt{1-|t|})}{(1-|t|)}.$$

Estimate for I_3 . We have

$$\begin{aligned} I_3 &= |F_t(\tau)| |\ln |F_t(\tau)|| \int_{\Gamma_2} \frac{|d\gamma|}{|\gamma-t|^2} \lesssim |F_t(\tau)| |\ln |F_t(\tau)|| \int_{\omega(u) \geq |F_t(\tau)|} \frac{du}{u^2 + (1-|t|)^2} \\ &= |F_t(\tau)| |\ln |F_t(\tau)|| \lesssim \frac{|F_t(\tau)| |\ln |F_t(\tau)||}{(1-|t|)} \left(\frac{\pi}{2} - \arg \cot \frac{x^*}{1-|t|} \right), \end{aligned} \tag{11}$$

where x^* is introduced by identity (10).

Taking into consideration the elementary inequality

$$0 \leq \frac{\pi}{2} - \arg \cot V \leq \frac{H}{1+V}, \quad V \in [0, +\infty),$$

by estimate (11) we finally obtain

$$I_3 \lesssim |F_t(\tau)| |\ln |F_t(\tau)|| \frac{H}{1-|t| + x^*}.$$

Now, in view of the estimates

$$\sup_{0 \leq u \leq 2} u |\ln u| \leq e, \quad x^* \geq \sqrt{1-|t|},$$

by (11) we get

$$I_3 \lesssim \frac{1}{\sqrt{1-|t|}} \frac{\omega(\sqrt{1-|t|})}{\sqrt{1-|t|}} = C_f \frac{\omega(\sqrt{1-|t|})}{1-|t|}.$$

In the latter inequality we employed the inequality $\frac{\omega(\delta)}{\delta} \geq \omega(1)$ as $0 < \delta \leq 1$.

2°. *Estimate for $|F_t(\tau)| |f'_t(t)|$* As above, we let

$$K_t(\xi) = \frac{1}{(t-\xi)^2}, \quad \xi \in \Gamma, \quad t \in D.$$

Then

$$f'_t(t) = f_t(t) \int_{\Gamma} K_t(\xi) d\mu^t(\xi),$$

where the measure μ^t is supported in the set

$$E_t = \{ \gamma \in \Gamma : |f(\gamma)| \leq \omega(\sqrt{1-|t|}) \},$$

at that, $\mu^t(\Gamma) \leq M$.

Let $\tau^* \in E_t$ be the point closest to the point t . Then by Lemma 2 we get

$$|F_t(t)| \lesssim \left[|F_t(\tau)| + \omega(1-|t|) \ln \frac{1}{1-|t|} \right]. \tag{12}$$

Hence,

$$|F_t(t)| \lesssim \left(|F_t(\tau) - F_t(\tau^*)| + |F_t(\tau^*)| \omega(1 - |t|) \ln \frac{1}{1 - |t|} \right).$$

Since $\tau^* \in E_t$, then $|F_t(\tau^*)| \leq \omega(\sqrt{1 - |t|})$. Therefore, by estimate (12) we obtain

$$|F_t(t)| \lesssim \left[\omega(|\tau - \tau^*|) + \omega(1 - |t|) \ln \frac{1}{1 - |t|} + \omega(\sqrt{1 - |t|}) \right].$$

By Lemma 1 we have

$$|F_t(t)| |f'_t(t)| \lesssim \left[\omega(|\tau - \tau^*|) + \omega(\sqrt{1 - |t|}) \right] |f_t(t)| \int_{\Gamma} |K_t(\xi)| d\mu^t(\xi),$$

that is,

$$|F_t(t)| |f'_t(t)| \lesssim \left[|f'_t(t)| \omega(|\tau - \tau^*|) + |f'_t(t)| \omega(\sqrt{1 - |t|}) \right].$$

We proceed to estimating the expression in the brackets.

Let

$$J_1 = |f'_t(t)| \omega(|\tau - \tau^*|) = \omega(|\tau - \tau^*|) |f_t(t)| \int_{\Gamma} \frac{d\mu^t(\xi)}{|\xi - t|^2},$$

$$J_2 = |f'_t(t)| \omega(\sqrt{1 - |t|}).$$

We first estimate J_2 .

We have

$$\begin{aligned} J_2 &\lesssim \omega(\sqrt{1 - |t|}) \int_{\Gamma} \frac{d\mu^t(\xi)}{|\xi - t|^2} \exp\left(-\frac{1 - |t|^2}{|\xi - t|^2} d\mu^t(\xi)\right) \\ &\lesssim \frac{\omega(\sqrt{1 - |t|})}{1 - |t|} \sup_{u>0} e^{-u} u \lesssim \frac{\omega(\sqrt{1 - |t|})}{1 - |t|}. \end{aligned} \quad (13)$$

We proceed to estimating J_1 . If $\omega(|\tau - \tau^*|) \lesssim \omega(\sqrt{1 - |t|})$, then J_1 can be estimated exactly in the same way as J_2 . This is why we assume that $\omega(\sqrt{1 - |t|}) \leq \omega(|\tau - \tau^*|)$. In view of the monotonicity of the function ω , the latter estimate implies $\sqrt{1 - |t|} \leq |\tau - \tau^*|$. Therefore, we obtain

$$\begin{aligned} J_1 &\lesssim \omega(|\tau - \tau^*|) |f_t(t)| \frac{1}{|t - \tau^*|^2} \lesssim \frac{\omega(|\tau - \tau^*|)}{|t - \tau^*|} \frac{1}{|t - \tau^*|} \\ &\lesssim \frac{\omega(|\tau - \tau^*|)}{|\tau - \tau^*|} \frac{1}{\sqrt{1 - |t|}} \lesssim \frac{\omega(\sqrt{1 - |t|})}{(\sqrt{1 - |t|})(\sqrt{1 - |t|})} = \frac{\omega(\sqrt{1 - |t|})}{1 - |t|}, \quad t \in D. \end{aligned} \quad (14)$$

In the latter inequality we have employed the non-increasing of the function $\frac{\omega(\delta)}{\delta}$ on $(0, 2)$.

Estimates (13), (14) imply the statement of the theorem. \square

Let us outline the proof of Theorem 2.

Let f satisfy the assumptions of Theorem 2. We consider the following cut-off function: $f(\lambda) = f(\lambda\xi)$, $\lambda \in D$, $\xi \in S_n$, a point ξ is fixed, see [11].

It is easy to see that $f_\xi(\lambda)$ satisfies the assumptions of Theorem 1. In view of Remark 2, we establish the estimate

$$|f_\xi(\lambda_1) - f_\xi(\lambda_2)| \leq A\omega\left(\sqrt{|\lambda_1 - \lambda_2|}\right), \quad \lambda_1, \lambda_2 \in D,$$

and A is independent of $\xi \in S_n$.

Employing the identity

$$R(f)(z) = n \int_{S_n} d\sigma(\xi) \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{\langle z, \xi \rangle e^{-i\theta} f(e^{i\theta} \xi) d\theta}{(1 - \langle z, \xi \rangle e^{-i\theta})^{n+1}}, \quad z \in B_n,$$

where R is the radial derivative, see [11], we get the estimate

$$|R(f)(z)| \lesssim \frac{\omega(\sqrt{1 - \|z\|})}{1 - \|z\|}, \quad z \in B_n.$$

Arguing as in the proof of Theorem 7.9 in [11], we arrive at the statement of Theorem 2.

BIBLIOGRAPHY

1. N.K. Bari, S.B. Stechkin. *Best approximations and differential properties of two conjugate functions* // Trudy Mosk. Matem. Obsch. **5**, 482–522 (1956). (in Russian).
2. A.V. Vasin, S.V. Kislyakov, A.N. Medvedev. *Local smoothness of an analytic function compared to the smoothness of its modulus* // Alg. Anal. **25:3**, 52–85 (2013). [St. Petersburg Math. J. **25:3**, 397–420 (2013).]
3. Ya.L. Geronimus. *On some properties of analytic functions continuous on a closed circle or circular sector* // Matem. Sborn/ **38(80):3**, 319–330 (1956). (in Russian).
4. N.P. Kornejchuk. *Extremal problems of approximation theory*. Nauka, Moscow (1976). (in Russian).
5. V.P. Havin, F.A. Shamoyan. *Analytic functions with the boundary values having Lipschitz module* // Zapis. Nauchn. Semin. LOMI. **19**, 237–239 (1970). (in Russian).
6. V.P. Khavin. *A generalization of the Privalov-Zygmund theorem on the modulus of continuity of the conjugate function* // Izv. Akad. Nauk Arm. SSSR. Ser. Matem. **6**, 252–258, 265–287 (1971).
7. F.A. Shamoyan. *Some division problems in spaces of analytic functions*. PhD thesis, Leningrad State Univ., Leningrad (1970). (in Russian).
8. N.A. Shirokov. *Outer functions from the analytic O.V. Besov classes* // Zapis. Nauch. Semin. POMI. **217**, 172–217 (1994). [J. Math. Sci. **85:2**, 1867–1897 (1997).]
9. N.A. Shirokov. *Smoothness of a holomorphic function in a ball and smoothness of its modulus on the sphere* // Zapis. Nauch. Semin. POMI. **447**, 123–127 (2016). (in Russian).
10. N.A. Shirokov. *Analytic functions smooth up to the boundary*. Lect. Notes Math. **1312**. Springer, Berlin (1988).
11. Kehe Zhu. *Space of holomorphic functions in the unit ball*. Graduate Texts Math. **226**. Springer, Berlin (2005).

Faizo Agitovich Shamoyan,
 Bryansk State University
 named after Academician Ivan Georgiyevich Petrovsky
 Bezhitsckaya str. 14,
 241036, Bryansk, Russia
 E-mail: shamoyanfa@yandex.ru