

# CONTINUABILITY OF MULTIPLE POWER SERIES INTO SECTORIAL DOMAIN BY MEANS OF INTERPOLATION OF COEFFICIENTS

A.J. MKRTCHYAN

**Abstract.** We consider the problem on continuability of a multiple power series centered at the origin of  $\mathbb{C}^n$  into a sectorial domain. The condition of the mentioned continuability is given in terms of an entire function interpolating the coefficients of the power series. We estimate the indicator function of the interpolating function with the help of which the sectorial set is determined. More precisely, the growth of the interpolating function on the imaginary subspace describes the sectorial set on which the series sum is continued. In the study we use methods of multivariate complex analysis, in particular, integral representations (Cauchy, Mellin, and Lindelof representations), multidimensional residues and properties of power series.

**Keywords:** multiple power series, analytic continuation, indicator of entire function, multidimensional residues.

**Mathematics Subject Classification:** 32A05, 30B30

## 1. INTRODUCTION

Analytic functions play a very important role in mathematics. One way to study an analytic function is to expand it into a power series. The coefficients of such power series contain the information of its analytic continuation. The problem on analytic continuation and a closely related problem of relationship between singularities of power series and its coefficients were extensively studied by Hadamard, Lindelöf, Pólya, Szegő, Carlson and many other prominent mathematicians, see the references in monograph by Biberbach [1]. One possible approach to treat this problem is to interpolate the coefficients by values  $\varphi(k)$  of an entire function  $\varphi(z)$  at a natural numbers  $k \in \mathbb{N}$ .

In case of a one variable power series, Arakelian [2], [3] gave a criterion for a given arc of a unit circle to be an arc of regularity in terms of the indicator function of the interpolating entire function. Pólya found conditions for analytic continuability of a series into the entire complex plane except for some boundary arc [4].

In the multivariable case, for some classes of power series, in [5] a condition for continuability into a sectorial domain was given in terms of a holomorphic function that interpolated the series coefficients.

---

A.J. MKRTCHYAN, CONTINUABILITY OF MULTIPLE POWER SERIES INTO SECTORIAL DOMAIN BY MEANS OF INTERPOLATION OF COEFFICIENTS.

© MKRTCHYAN A.J. 2022.

THE RESEARCH IS SUPPORTED BY GRANT OF THE RUSSIAN SCIENCE FOUNDATION (PROJECT NO. 20-11-20117).

*Submitted September 15, 2021.*

We consider the multiple power series

$$f(z) = \sum_{k \in \mathbb{Z}_+^n} f_k z^k \tag{1.1}$$

with the property

$$\overline{\lim}_{|k| \rightarrow \infty} |k| \sqrt{|f_k| R^k} = 1, \tag{1.2}$$

where  $R^k = R_1^{k_1} \dots R_n^{k_n}$  and  $|k| = k_1 + \dots + k_n$ . According to the  $n$ -dimensional Cauchy-Hadamard theorem [6, Sect. 7], property (1.2) means that  $R_j$  constitute the family of radii of polydisk of convergence of series (1.1). In this work, we give sufficient conditions for analytic continuability of a multiple power series into a sectorial domain. A domain  $G \subset \mathbb{C}^n$  is called sectorial if it is defined by the conditions on the arguments  $\theta = (\arg z_1, \dots, \arg z_n)$  of elements  $z \in \mathbb{C}^n$  only.

For  $n$  variables, by an entire function of exponential type we mean a function  $\varphi(z) = \varphi(z_1, \dots, z_n)$  holomorphic in  $\mathbb{C}^n$ , for which there exist positive  $A, \sigma_1, \dots, \sigma_n$  such that for all  $z \in \mathbb{C}^n$  an inequality

$$|\varphi(z)| \leq A e^{\sigma_1 |z_1| + \dots + \sigma_n |z_n|}$$

holds.

Following Ivanov [7], we introduce the set, which implicitly contains the notion of the growth indicator of an entire function of exponential type  $\varphi(z) \in \mathcal{O}(\mathbb{C}^n)$ :

$$T_\varphi(\theta) = \{ \nu \in \mathbb{R}^n : \ln |\varphi(r e^{i\theta})| \leq \nu_1 r_1 + \dots + \nu_n r_n + C_{\nu, \theta} \},$$

where the inequality is satisfied for each  $r \in \mathbb{R}_+^n$  with some constant  $C_{\nu, \theta}$ . Here  $r e^{i\theta}$  stands for the vector  $(r_1 e^{i\theta_1}, \dots, r_n e^{i\theta_n})$ . Thus,  $T_\varphi(\theta)$  is the set of linear majorants (up to a shift by  $C_{\nu, \theta}$ )

$$\nu(r) = \nu_1 r_1 + \dots + \nu_n r_n$$

for the logarithm of the absolute value of the function  $\varphi$ .

We denote

$$T_\varphi := \bigcap_{\theta_j = \pm \frac{\pi}{2}} T_\varphi(\theta_1, \dots, \theta_n),$$

$$\mathcal{M}_\varphi := \{ \nu \in [0, \pi)^n : \nu + \varepsilon \in T_\varphi, \nu - \varepsilon \notin T_\varphi \text{ for any } \varepsilon \in \mathbb{R}_+^n \}.$$

Let  $G$  be a sectorial set

$$G = \bigcup_{\nu \in \mathcal{M}_\varphi} \{ z \in \mathbb{C}^n : \nu_j < \arg z_j < 2\pi - \nu_j, j = 1, \dots, n \}. \tag{1.3}$$

This set is a domain: it is open and connected because every polysector  $G_\nu$  is connected and contains the point  $(-1, \dots, -1)$ .

**Theorem 1.** *Let  $\varphi(\zeta)$  be an entire function of the exponential type interpolating coefficients  $f_k$  of series (1.1). Assume that there exists  $\nu(\theta) \in \overline{T}_\varphi(\theta)$  for  $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]^n$  such that inequalities*

$$\nu_j(\theta) \leq a |\sin \theta_j| + b \cos \theta_j, \quad j = 1, \dots, n, \tag{1.4}$$

*hold with  $a \in [0, \pi)$ ,  $b \in [0, \infty)$ . Then the sum of the series can be continued analytically to a sectorial domain  $G$  of the form (1.3).*

## 2. PROOF OF THEOREM

Let  $\varphi$  be an entire function obeying the assumptions of the theorem. For each  $\nu \in T_\varphi(\theta)$ , this function satisfies the inequality

$$|\varphi(re^{i\theta})| \leq A_{\nu,\theta} e^{(\nu,r)} \quad \forall r \in \mathbb{R}_+^n.$$

Then for  $\nu \in \bar{T}_\varphi(\theta)$  (closure  $T$ ) it satisfies the inequality

$$|\varphi(re^{i\theta})| \leq A_{\nu,\theta} e^{(\nu,r)+o(r)} \quad \text{for all } r \in \mathbb{R}_+^n.$$

By (1.4) we find that for  $|\theta_j| \leq \frac{\pi}{2}$ ,  $j = 1, \dots, n$ ,

$$|\varphi(re^{i\theta})| \leq A_{\nu,\theta} e^{a \sum_{j=1}^n r_j |\sin \theta_j| + b \sum_{j=1}^n r_j \cos \theta_j + o(r)}.$$

Representing

$$\zeta_j = \xi_j + i\eta_j = r_j e^{i\theta_j},$$

we get

$$|\varphi(\zeta)| \leq A_{\nu,\theta} e^{a \sum |\eta_j| + b \sum \xi_j + o(|\zeta|)} \quad (2.1)$$

for  $\zeta_j \in \Delta_{\pi/2}$ ,  $j = 1, \dots, n$ .

We consider the following function:

$$g(\zeta, z) = \prod_{j=1}^n \frac{z_j^{\zeta_j}}{(e^{2\pi i \zeta_j} - 1)},$$

where  $z_j = x_j + iy_j$ ,  $j = 1, \dots, n$ . It is meromorphic in  $\zeta \in \mathbb{C}^n$  and holomorphic in  $z \in (\mathbb{C} \setminus \mathbb{R}_+)^n$ .

Let  $D_\rho(a) := \{z \in \mathbb{C} : |z - a| < \rho\}$  be an open circle with center  $a \in \mathbb{C}$  and radius  $\rho > 0$ . We let

$$D^* := \bigcup_{m \in \mathbb{Z}} D_{\delta/2}(m),$$

where  $\delta < e^{-b}$  and we observe that there exists a constant  $C = C_\delta > 0$  such that

$$|e^{2\pi i w} - 1| > \frac{e^{\pi(|\operatorname{Im} w| - \operatorname{Im} w)}}{C} \quad \text{for } w \in D^*.$$

Therefore, we have the estimate:

$$g(\zeta, z) < C e^{(\xi, \log |z|) - \langle (\pi - |\pi - \arg z|), |\eta| \rangle} \quad (2.2)$$

for  $\zeta \in (\mathbb{C} \setminus D^*)^n$  and  $z \in (\mathbb{C} \setminus \mathbb{R}_+)^n$ .

Using (2.1) and (2.2) for  $\zeta \in (\Delta_{\pi/2} \setminus D^*)^n$  and  $z \in (\mathbb{C} \setminus \mathbb{R}_+)^n$ , we obtain

$$|\varphi(\zeta)| |g(\zeta, z)| \leq c e^{\sum \xi_j (\ln |z_j| + b) - \sum (\pi - a - |\pi - \arg z_j|) |\eta_j| + o(|\zeta|)}.$$

Let

$$K = \bar{D}_{e^{-b-\delta}} \setminus (\Delta_{a+\delta}^o \cup D_\delta).$$

We note that

$$\pi - a - |\pi - \arg z_j| \geq \delta \quad \text{as } z_j \in K, \quad j = 1, \dots, n.$$

Hence, for  $z \in K^n$  and  $\zeta \in (\Delta_{\pi/2} \setminus D^*)^n$  we get

$$|g(\zeta, z)| |\varphi(\zeta)| < c_1 e^{\sum \xi_j (\ln |z_j| + b) - \delta \sum |\eta_j| + o(|\zeta|)}. \quad (2.3)$$

For each  $m = (m_1, \dots, m_n) \in \mathbb{N}^n$  we consider the integral

$$I_m = \int_{\partial G_m} \varphi(\zeta) g(\zeta, z) d\zeta, \quad (2.4)$$



Taking into consideration (2.5), we see that

$$I_m = \sum_{k_1=1}^{m_1} \dots \sum_{k_n=1}^{m_n} \varphi(k_1, \dots, k_n) z_1^{k_1} \dots z_n^{k_n}.$$

We consider the integral

$$\mathbb{I} = \int_{\Gamma^1} g(\zeta, z) \varphi(\zeta) d\zeta, \quad \Gamma^1 = \Gamma_1^1 \times \dots \times \Gamma_1^n,$$

where

$$\Gamma_j^1 = [-i\infty, -i\delta] \cup \left\{ \delta e^{i\theta_j}, |\theta_j| \leq \frac{\pi}{2} \right\} \cup [i\delta, i\infty], \quad j = 1, \dots, n.$$

We are going to show that for  $\zeta \in \Gamma^1$  and  $z \in G$ , the absolute value of the integrand  $|\varphi(\zeta)||g(\zeta, z)|$  is estimated by

$$|g(\zeta, z)||\varphi(\zeta)| < c(z)e^{-\delta \sum |\eta_j|}.$$

Indeed, it follows from the definition of the set  $T_\varphi$  ( $T_\varphi$  describes the growth of the function  $\varphi$  along the imaginary subspace) that

$$|\varphi(\zeta)| \leq e^{C_{\nu, \theta}} e^{\sum \nu_j |\eta_j|} \quad \text{as } \zeta \in \Gamma^1,$$

where  $\nu_j$  is a  $j$ th component of the vector  $\nu$ , which ranges in the set  $T_\varphi$ . By the previous inequality and (2.2), for  $\zeta \in \Gamma^1$  and  $z \in (\mathbb{C} \setminus \mathbb{R}_+)^n$  we get

$$|\varphi(\zeta)||g(\zeta, z)| \leq c(|z|)e^{-\sum \pi - \nu_j - |\pi - \arg z_j| |\eta_j|}.$$

We observe that

$$\pi - \nu_j - |\pi - \arg z_j| \geq \delta$$

for  $z \in (\mathbb{C} \setminus \Delta_{\nu_1 + \delta}^o) \times \dots \times (\mathbb{C} \setminus \Delta_{\nu_n + \delta}^o)$  for each  $\nu_j < \pi$ . By two latter inequalities and Proposition 1 we have

$$|\varphi(\zeta)||g(\zeta, z)| \leq C e^{-\delta \sum |\eta_j|}$$

for  $\zeta \in \Gamma^1$  and

$$z \in \bigcup_{\{\nu \in T_\varphi, \nu_j < \pi\}} G_{\nu + \delta}. \tag{2.6}$$

It follows from the definition of  $\mathcal{M}_\varphi$  that (2.6) is equivalent to

$$z \in \bigcup_{\nu \in \mathcal{M}_\varphi} G_\nu.$$

Hence, the integral  $\mathbb{I}$  converges for  $z \in G$ . Since  $I_m \rightarrow \mathbb{I}$ , as  $m_j \rightarrow \infty$ ,  $j = 1, \dots, n$ , for  $z \in (K^o)^n$  we get

$$\mathbb{I}(z) = f(z) - \sum_{j=1}^n f_j(z) \quad \text{where} \quad f_j = \sum_{k_1 \dots k_n} f_{k_1 \dots k_n} z_1^{k_1} \dots z_n^{k_n}.$$

We observe that the functions  $\varphi_j = \varphi(z_1, \dots, z_n)$  are interpolation functions for  $(n - 1)$ -dimensional series  $f_j(z)$  and they satisfy the assumptions of the theorem. Moreover, the set  $T_{\varphi_j}$  contains the subset  $T_\varphi$  without the component  $\nu_j$ . This means that the series  $f_j(z)$  in  $\mathbb{C}$  can be continued into a sectorial sets, which contains  $G$ . Therefore, the sum of series (1.1) can be continued analytically into the sectorial set  $G$  and this completes the proof.

## 3. EXAMPLE

We consider a power series

$$f(z_1, z_2) = \sum_{k_1, k_2 \in \mathbb{N}^2} \cos \sqrt{k_1 k_2} z_1^{k_1} z_2^{k_2}. \quad (3.1)$$

It obvious that the function  $\varphi(\zeta_1, \zeta_2) = \cos(\zeta_1 \zeta_2)^{\frac{1}{2}}$  is entire and it interpolates the coefficients of series (3.1). Representing  $\zeta_j = r_j e^{i\theta_j}$ , for the absolute value of the function we get an asymptotic expansion

$$|\varphi(\zeta_1, \zeta_2)| = \left| \cos \left( (r_1 r_2)^{\frac{1}{2}} e^{i \frac{\theta_1 + \theta_2}{2}} \right) \right| = \frac{1}{2} e^{(r_1 r_2)^{\frac{1}{2}} \left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right|} + o(1)$$

as  $r_1 r_2 \rightarrow \infty$ . Therefore, the set  $T_\varphi(\theta)$  reads as

$$T_\varphi(\theta) = \left\{ \nu \in \mathbb{R}^2 : (r_1 r_2)^{\frac{1}{2}} \left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right| \leq \nu_1 r_1 + \nu_2 r_2 + C_{\nu, \theta} \right\},$$

and, hence, it consists of solutions  $\nu = \nu(\theta)$  of the inequality

$$(r_1 r_2)^{\frac{1}{2}} \left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right| \leq \nu_1 r_1 + \nu_2 r_2, \quad r_1, r_2 \geq 0.$$

Taking  $r_j = 0$ , we see that  $\nu_j \geq 0$ ,  $j = 1, 2$ .

In order to study this inequality for  $r_1 r_2 \neq 0$ , we take into consideration that it is homogeneous with respect to  $r_1$  and  $r_2$ . Namely, dividing it by  $r_2$ , we find:

$$\left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right| \leq \nu_1 \left( \frac{r_1}{r_2} \right)^{\frac{1}{2}} + \nu_2 \left( \frac{r_2}{r_1} \right)^{\frac{1}{2}};$$

we then denote

$$t = \left( \frac{r_1}{r_2} \right)^{\frac{1}{2}}.$$

Then the inequality is reduced to the following inhomogeneous inequality

$$\nu_1 t^2 - \left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right| t + \nu_2 \geq 0, \quad t \geq 0. \quad (3.2)$$

As it has been stated above, we are interested only in solutions  $\nu$  with nonnegative coordinates. It follows from Viéte's formulas that for  $\nu_1 \geq 0, \nu_2 \geq 0$  the quadratic trinomial (3.2) in  $t$  has no negative roots, therefore we may consider the inequality for all  $t \in \mathbb{R}$ . Thus, the solutions  $\nu$  of inequality (3.2) are defined by the condition that its discriminant is non-positive:

$$\left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right|^2 - 4\nu_1 \nu_2 \leq 0,$$

that is,

$$\nu_1 \nu_2 \geq \frac{1}{4} \left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right|^2.$$

Finally, we get

$$\bar{T}_\varphi(\theta_1, \theta_2) = \left\{ \nu \in \mathbb{R}^2 : \nu_1 \nu_2 = \frac{1}{4} \left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right|^2, \quad \nu_1 \geq 0, \quad \nu_2 \geq 0 \right\}. \quad (3.3)$$


Obviously,  $T_\varphi(\pm \frac{\pi}{2}, \pm \frac{\pi}{2})$  consists of the quadrant  $\nu_1 \geq 0, \nu_2 \geq 0$  if the signs are the same, and  $\{\nu \in \mathbb{R}_+^2 : \nu_1 \nu_2 \geq \frac{1}{4}\}$  if the signs are different.

As a result, the intersection  $T_\varphi = \bigcap T_\varphi(\pm\frac{\pi}{2}, \pm\frac{\pi}{2})$  is

$$T_\varphi = \left\{ \nu \in \mathbb{R}^2 : \nu_1 \nu_2 \geq \frac{1}{4}, \nu_1 \geq 0, \nu_2 \geq 0 \right\}.$$

Therefore, the sets  $\mathcal{M}_\varphi$  read as

$$\mathcal{M}_\varphi = \left\{ \nu \in [0, \pi)^2 : \nu_1 \nu_2 = \frac{1}{4} \right\}.$$



Now let us confirm that condition (1.4) of Theorem is satisfied. In view of (3.3), it is sufficient to find constants  $a \in [0, \pi)$ ,  $b \in [0, \infty)$  such that


$$(a|\sin \theta_1| + b \cos \theta_1)(a|\sin \theta_2| + b \cos \theta_2) \geq \frac{1}{4} \left| \sin \left( \frac{\theta_1 + \theta_2}{2} \right) \right|^2.$$

For  $a = 1, b = 1$  the left hand side satisfies

$$(|\sin \theta_1| + \cos \theta_1)(|\sin \theta_2| + \cos \theta_2) \geq 1,$$

and the right hand side does not exceed  $\frac{1}{4}$  and this proves the inequality.

Hence, according to Theorem, the sum of the series can be continued into the union  $G$  of the polysectors  $G_\nu = (\mathbb{C} \setminus \Delta_{\nu_1}) \times \dots \times (\mathbb{C} \setminus \Delta_{\nu_n})$  over all  $\nu \in \mathcal{M}_\varphi$ . The Figure below depicts the set of the arguments  $\theta = (\theta_1, \theta_2)$  defining the sectorial set  $G$ . It is a union of rectangles  $(\nu_1, 2\pi - \nu_1) \times (\nu_2, 2\pi - \nu_2)$  over all  $(\nu_1, \nu_2) \in \mathcal{M}_\varphi$ .



## REFERENCES

1. L. Bieberbach. *Analytische Fortsetzung*. Springer-Verlag, Berlin (1955).
2. N.U. Arakelian. *Approximation by entire functions and analytic continuation* // in “Progress in approximation theory. An international perspective”, Proc. International Conference on approximation theory, Tampa, South Florida, USA, 1990. Springer-Verlag, New York, 295–313 (1992).
3. N. Arakelian, W. Luh, J. Muller. *On the localization of singularities of lacunar power series* // Compl. Var. Ellip. Equats. **52**:7, 561–573 (2007).
4. G. Pólya. *Untersuchungen über Lücken und Singularitäten von Potenzreihen* // Math. Zeit. **29**, 549–640 (1929).
5. A.J. Mkrtchyan. *Continuation of multiple power series in a sectorial domain* // Math. Nach. **292**:11, 2441–2451 (2019).
6. B.V. Shabat. *Introduction to Complex Analysis: functions of several variables*. Nauka, Moscow (1976). [Am. Math. Soc., Providence, RI (1992).]
7. V.K. Ivanov. *A characterization of the growth of an entire function of two variables and its application to the summation of double power series* // Matem. Sb. **47(89)**:1, 3–16 (1959). (in Russian).
8. L.I. Ronkin. *Introduction to the theory of entire functions of several variables*. Nauka, Moscow (1971). [Am. Math. Soc., Providence, RI (1974).]
9. A.K. Tsikh. *Multidimensional residues and their applications*. Nauka, Novosibirsk (1988). [Am. Math. Soc., Providence, (1992).]
10. O. N. Zhdanov, A. K. Tsikh. *Investigation of multiple Mellin-Barnes integrals by means of multidimensional residues* // Siber. Math. J. **39**:2, 281–298 (1998).

Aleksandr Janibekovich Mkrtchyan,  
Institute of Mathematics and Computer Science, Siberian Federal University,  
Svobodny 79,  
660041, Krasnoyarsk, Russia  
Institute of Mathematics of National Academy of Republic Armenia,  
Marshal Baghramian 24/5,  
0019, Yerevan, Armenia  
E-mail: Alex0708@bk.ru