

INVERTING OF GENERALIZED RIEMANN-LIOUVILLE OPERATOR BY MEANS OF INTEGRAL LAPLACE TRANSFORM

I.I. BAVRIN, O.E. YAREMKO

Abstract. We employ the integral Laplace transform to invert the generalized Riemann-Liouville operator in a closed form. We establish that the inverse generalized Riemann-Liouville operator is a differential or integral-differential operator. We establish a relation between Riemann-Liouville operator and Temlyakov-Bavrin operator. We provide new examples of generalized Riemann-Liouville operator.

Keywords: Riemann-Liouville operator, fractional integral, Laplace transform.

Mathematics Subject Classification: 26A33, 44A10

1. INTRODUCTION

Let $f(x) \in L_1(0, 1)$, then the function

$$J^\alpha [f](x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt \in L_1(0, 1)$$

is called a fractional integral of order α , see [8, 11, 15]. The fractional integral can be written as

$$J^\alpha [f](x) = \frac{x^\alpha}{\Gamma(\alpha)} \int_0^1 (1-\varepsilon)^{\alpha-1} f(\varepsilon x) d\varepsilon \in L_1(0, 1).$$

The latter formula allows one to extend the fractional integration for the case of the functions $w = f(z)$ analytic in the unit circle $B = \{z : |z| < 1\}$. As a result, we arrive at the generalized Riemann-Liouville operator

$$L_\omega [f](z) = \alpha \int_0^1 (1-\varepsilon)^{\alpha-1} f(\varepsilon z) d\varepsilon. \quad (1.1)$$

The extension of the notion of fractional integral (1.1) was made by M.M. Džrbašjan. In the work [5], the generalized Riemann-Liouville operator was introduced.

We shall say a function $\omega(x)$ belongs to Ω if it is nonnegative and continuous on $[0, 1)$ and

$$\omega(0) = 1, \int_0^1 \omega(x) dx < \infty,$$

for all $r \in [0, 1)$ the inequality

$$\int_r^1 \omega(x) dx > 0$$

holds true. Let $w = f(z)$ be a function analytic in the unit circle $B = \{z : |z| < 1\}$.

I.I. BAVRIN, O.E. YAREMKO, INVERTING OF GENERALIZED RIEMANN-LIOUVILLE OPERATOR BY MEANS OF INTEGRAL LAPLACE TRANSFORM.

© BAVRIN I.I., YAREMKO O.E. 2016.

Definition 1.1. *The generalized Riemann-Liouville operator is the following operator [5]*

$$L_\omega [f] = f(0) + z \int_0^1 \omega(\varepsilon) f'(\varepsilon z) d\varepsilon.$$

We define the sequence of the numbers

$$\Delta_1 = 1, \Delta(k) = k \int_0^1 r^{k-1} \omega(x) dx < \infty, \quad k = 1, 2, \dots$$

M.M. Džrbašjan established the following statements [5]:

i)

$$L_\omega [z^k] = \Delta_k z^k;$$

ii) Let $w = f(z)$ be a function analytic in the unit circle and assume that its Taylor series is of the form

$$f(z) = \sum_{k=0}^{\infty} a_k z^k.$$

Then

$$L_\omega [f](z) = \sum_{k=0}^{\infty} a_k \Delta_k z^k;$$

iii) The operator L_ω is invertible and

$$L_\omega^{-1} [f](z) = \sum_{k=0}^{\infty} a_k \frac{z^k}{\Delta_k},$$

where

$$f(z) = \sum_{k=0}^{\infty} a_k z^k.$$

iiii) If a function $\omega(x) \in \Omega$ is a continuously differentiable in $[0, 1)$ and $\omega(1) = 0$, then the generalized Riemann-Liouville operator is of the form

$$L_\omega [f] = - \int_0^1 \omega'(\varepsilon) f(\varepsilon z) d\varepsilon. \quad (1.2)$$

The current state-of-art of the fractal analysis and its applications is presented in [15-18].

2. MAIN RESULT

As it was mentioned, the theory of the generalized Riemann-Liouville operator was constructed by M.M. Džrbašjan in work [5]. However, no explicit construction for the inverse operator was proposed. In order to invert the generalized Riemann-Liouville operator in a closed form, we propose to employ the integral Laplace transform.

The idea is to continue the scalar sequence

$$\Delta(k) = k \int_0^1 r^{k-1} \omega(x) dx < \infty, k = 1, 2, \dots$$

into the half-plane $p: \operatorname{Re} p = \sigma > \sigma_0 \geq 0$.

The problem on inverting the generalized Riemann-Liouville operator is solved if the function $\Delta(p)$ is a Laplace image, and the function

$$\frac{1}{\Delta(p)}$$

has a power growth, i.e.,

$$\frac{1}{\Delta(p)} = p^\alpha L(p), \alpha > 0,$$

where $L(p)$ is a slowly varying function [14].

Lemma 2.1. *The function*

$$\Delta(p) = p \int_0^1 \varepsilon^{p-1} \omega(\varepsilon) d\varepsilon$$

is well-defined for the values $p \operatorname{Re} p = \sigma > \sigma_0 \geq 0$ and serves as the Laplace image of the function

$$\omega(e^{-t}), \omega(e^{-t}) \geq 0, t \in [0, \infty).$$

Proof. It is sufficient to make a change of the variable $\varepsilon = e^{-t}$ in the expression for $\Delta(p)$. \square

Theorem 2.1. *Suppose that all roots of the polynomial $Q(p)$ are located in the left half-plane and $Q(0) \neq 0$ and let the function*

$$\frac{1}{Q(p) \Delta(p)}$$

be the image of some original $\omega^*(t)$. The the inverse operator for L_ω is of the form

$$L_\omega^{-1}[f](z) = Q\left(z \frac{d}{dz}\right) \left[\int_0^\infty \omega^*(\varepsilon) f(e^{-\varepsilon} z) d\varepsilon \right], \quad (2.1)$$

where

$$Q\left(z \frac{d}{dz}\right) = a_0 + a_1 z \frac{d}{dz} + a_2 \left(z \frac{d}{dz}\right)^2 + \dots + a_n \left(z \frac{d}{dz}\right)^n,$$

numbers a_k are the coefficients of the polynomial $Q(p)$.

Proof. The integral component of the operator L_ω^{-1} that is, the operator of the form

$$\int_0^\infty \omega^*(\varepsilon) f(e^{-\varepsilon} z) d\varepsilon,$$

is continuous in the space of the functions $H(B) \cup C(\bar{B})$, and this is why

$$L_\omega^{-1}[f](z) = Q\left(z \frac{d}{dz}\right) \sum_{k=0}^{\infty} a_k z^k \frac{1}{Q(k) \Delta(k)} = \sum_{k=0}^{\infty} a_k z^k \frac{Q(k)}{Q(k) \Delta(k)} = \sum_{k=0}^{\infty} a_k z^k \frac{1}{\Delta(k)}.$$

This completes the proof. \square

Corollary 2.1. *The inverse for the Riemann-Liouville operator is either differential or integral-differential.*

Theorem 2.2. *Let the functions $\frac{1}{\Delta(p)} = Q(z)$ be a polynomial, then the inverse operator for L_ω is a differential operator of the form*

$$L_\omega^{-1}[f](z) = Q\left(z \frac{d}{dz}\right) [f(z)]. \quad (2.2)$$

Proof. We have the following identities

$$Q\left(z \frac{d}{dz}\right) \sum_{k=0}^{\infty} a_k z^k = \sum_{k=0}^{\infty} a_k z^k Q(k) = \sum_{k=0}^{\infty} a_k z^k \frac{1}{\Delta(k)} \equiv L_\omega^{-1}[f](z).$$

\square

3. VARIOUS CASES OF GENERALIZED RIEMANN-LIOUVILLE OPERATORS

3.1. Let $\omega(x) = 1 - x^h$, $h > 0$, then

$$L_\omega[f] = f(0) + z \int_0^1 (1 - \varepsilon^h) f'(\varepsilon z) d\varepsilon = h \int_0^1 \varepsilon^{h-1} f(\varepsilon z) d\varepsilon,$$

and we obtain the operator which differs by a factor from the operator

$$L_h^{-1}[f] = \int_0^1 \varepsilon^{h-1} f(\varepsilon z) d\varepsilon$$

studied in the works by I.I. Bavrin [1, 2]. In order to find the inverse operator we note that

$$\Delta(p) = p \int_0^1 \varepsilon^{p-1} (1 - \varepsilon^h) d\varepsilon = \frac{h}{h+p}.$$

Then, taking into consideration that the expression

$$\frac{1}{\Delta(p)} = \frac{h+p}{h}$$

is a polynomial, by Theorem 2.1 we obtain

$$L_\omega^{-1}[f] = \frac{hf + zf'(z)}{h} = \frac{L_h[f(z)]}{h}. \quad (3.1)$$

3.2. Let $\Delta(p) = \prod_{j=1}^m \frac{h_j}{h_j+p}$, $h_j > 0$. The original of the function $\Delta(p)$ is of the form

$$\sum_{k=1}^m e^{-h_k t} \prod_{j \neq k} \frac{h_j}{h_j - h_k} \rightarrow \Delta(p).$$

By formula (1.2) we establish the following expression for the operator L_ω

$$L_\omega[f](z) = \prod_{j \neq k} \frac{h_j}{h_j - h_k} \int_0^1 \varepsilon^{h_k-1} f(\varepsilon z) d\varepsilon.$$

At that, the inverse operator L_ω^{-1} is a differential operator of the form

$$L_\omega^{-1}[f](z) = \prod_{j=1}^m \frac{h_j + L_0}{h_j}.$$

The theory of such operators was exposed in the monographs by I.I. Bavrin [1, 2].

3.3. Let us discuss a fractional power of the operator $h + z \frac{d}{dz}$. We choose the function $\Delta(p)$ as

$$\Delta(p) = \frac{h^\alpha}{(h+p)^\alpha}.$$

Taking into consideration the formula

$$\frac{h^\alpha}{\Gamma(\alpha)} e^{-ht} t^{\alpha-1} \rightarrow \frac{h^\alpha}{(h+p)^\alpha}$$

in the table of Laplace transformations, according to formula (1.2), we establish the expression for the operator L_ω :

$$L_\omega[f](z) = \frac{h^\alpha}{\Gamma(\alpha)} \int_0^\infty e^{-h\varepsilon} \varepsilon^{\alpha-1} f(e^{-\varepsilon} z) d\varepsilon.$$

After the change of the variable $e^{-\varepsilon} = \tau$, we arrive at the following expression for the operator L_ω :

$$L_\omega[f](z) = \frac{h^\alpha}{\Gamma(\alpha)} \int_0^1 \varepsilon^{h-1} \left(\ln \frac{1}{\varepsilon} \right)^{\alpha-1} f(\varepsilon z) d\varepsilon, \quad h > 0, \quad \alpha > 0.$$

Thus, in the definition of the generalized Riemann-Liouville operator we should let

$$\omega(x) = \frac{h^\alpha}{\Gamma(\alpha)} \int_x^1 \varepsilon^{h-1} \left(\ln \frac{1}{\varepsilon} \right)^{\alpha-1} d\varepsilon.$$

a) If $\alpha = n \in N$, then the inverse operator L_ω^{-1} is calculated by the formula

$$L_\omega^{-1} = \left(h + z \frac{d}{dz} \right)^n,$$

i.e., is differential.

b) If $\alpha \notin N$, the inverse operator L_ω^{-1} is of the form:

$$\begin{aligned} L_\omega^{-1} [f] (z) &\equiv \left(h + z \frac{d}{dz} \right)^\alpha [f] (z) \\ &= \frac{1}{\Gamma(n+1-\alpha)} \left(h + z \frac{d}{dz} \right)^{n+1} \int_0^1 \varepsilon^{h-1} \left(\ln \frac{1}{\varepsilon} \right)^{n-\alpha} f(\varepsilon z) d\varepsilon. \end{aligned}$$

For instance, as $\alpha = \frac{1}{2}$, we obtain the following expression:

$$\left(h + z \frac{d}{dz} \right)^{\frac{1}{2}} [f] (z) = \frac{1}{\sqrt{\pi}} \left(h + z \frac{d}{dz} \right) \int_0^1 \varepsilon^{h-1} \left(\ln \frac{1}{\varepsilon} \right)^{-\frac{1}{2}} [f] (\varepsilon z) d\varepsilon.$$

3.4. Hadamard operator. We consider the class of functions $w = f(z)$ analytic in the unit circle $B = \{z : |z| < 1\}$ such that $f(0) = 0$. We choose the function $\Delta(p)$ as

$$\Delta(p) = \frac{1}{p^\alpha}.$$

Taking into consideration the formula

$$\frac{t^{\alpha-1}}{\Gamma(\alpha)} \rightarrow \frac{1}{p^\alpha}$$

from the table of the Laplace transformations, according to formula (2.1), we get the expression for the operator L_ω

$$L_\omega [f] (z) = \frac{1}{\Gamma(\alpha)} \int_0^\infty \varepsilon^{\alpha-1} f(e^{-\varepsilon} z) d\varepsilon, \quad \alpha > 0.$$

After the change of the variable $e^{-\varepsilon} = \tau$, we arrive at the following expression for operator L_ω

$$L_\omega [f] (z) = \frac{1}{\Gamma(\alpha)} \int_0^1 \left(\ln \frac{1}{\varepsilon} \right)^{\alpha-1} \frac{f(\varepsilon z)}{\varepsilon} d\varepsilon. \quad (3.2)$$

The right hand side of formula (3.2) defines Hadamard operator [12].

a) If $\alpha = n \in N$, the inverse operator L_ω^{-1} is calculated by the formula

$$L_\omega^{-1} = L_0^n,$$

i.e., is differential.

b) If $\alpha \notin N$, then inverse operator L_ω^{-1} is of the form:

$$L_\omega^{-1} [f] (z) \equiv L^\alpha [f] (z) = \frac{1}{\Gamma(n+1-\alpha)} L^{n+1} \left[\int_0^1 \left(\ln \frac{1}{\varepsilon} \right)^{n-\alpha} \frac{f(\varepsilon z)}{\varepsilon} d\varepsilon \right].$$

For instance, as $\alpha = \frac{1}{2}$, we obtain the following expression:

$$L^{\frac{1}{2}} [f] (z) = \frac{1}{\sqrt{\pi}} L \left[\int_0^1 \left(\ln \frac{1}{\varepsilon} \right)^{-\frac{1}{2}} \frac{f(\varepsilon z)}{\varepsilon} d\varepsilon \right].$$

3.5. If $\Delta(p) = \frac{h^2}{h^2+p^2}$, we obtain $Q(z) = \frac{h^2+z^2}{h^2}$, and, hence, the inverse operator is a differential operator of the form

$$L_\omega^{-1} = \frac{h^2 + L_0^2}{h^2}, L_0 = z \frac{d}{dz}.$$

Let us find the direct operator L_ω . By a formula in the table of the Laplace transformations,

$$h \sin ht \rightarrow \frac{h^2}{h^2 + p^2}.$$

According to Theorem 2.1, let us find the expression for the operator L_ω in formula (2.1)

$$L_\omega[f](z) = \int_0^\infty h \sin(h\varepsilon) f(e^{-\varepsilon}z) d\varepsilon.$$

As a result, after a change of the variable, we have

$$L_\omega[f](z) = \int_0^1 \frac{h}{\varepsilon} \sin\left(h \ln \frac{1}{\varepsilon}\right) f(\varepsilon z) d\varepsilon.$$

Remark 1. Here we have $\omega'(x) = -\frac{h}{x} \sin\left(h \ln \frac{1}{x}\right)$ and hence, $\omega(x) \notin \Omega$.

3.6. Fractional power of the operator $h^2 + L_0^2$.

We let

$$\Delta(p) = \frac{(h^2 + p^2)^\alpha}{h^{2\alpha}}, \quad h \neq 0, \quad \alpha > 0.$$

By a formula in the table of the Laplace transformations,

$$\frac{\sqrt{\pi} \cdot h}{\Gamma(\alpha)} \left(\frac{th}{2}\right)^{\alpha-\frac{1}{2}} J_{\alpha-\frac{1}{2}}(ht) \rightarrow \frac{h^{2\alpha}}{(h^2 + p^2)^\alpha},$$

where $J_{\alpha-\frac{1}{2}}(z)$ is the Bessel function of order $\alpha - \frac{1}{2}$. By Theorem 2.1 and formula (2.1) we find the expression for operator L_ω

$$L_\omega[f](z) = \frac{\sqrt{\pi}}{\Gamma(\alpha)} \int_0^1 \frac{h}{\varepsilon} \left(\frac{h \ln \frac{1}{\varepsilon}}{2}\right)^{\alpha-\frac{1}{2}} J_{\alpha-\frac{1}{2}}\left(h \ln \frac{1}{\varepsilon}\right) f(\varepsilon z) d\varepsilon.$$

a) If $\alpha = n \in N$, then the inverse Riemann-Liouville operator is a differential operator of the form

$$L_\omega^{-1} = \frac{(h^2 + L_0^2)^n}{h^{2n}}.$$

b) If $\alpha \notin N$, then the inverse operator L_ω^{-1} is of the form:

$$\begin{aligned} L_\omega^{-1}[f](z) &\equiv (h^2 + L_0^2)^\alpha [f](z) / h^{2\alpha} \\ &= \frac{\sqrt{\pi} h^{-2\alpha}}{\Gamma(n+1-\alpha)} \int_0^1 \varepsilon^{-1} \left(\frac{\ln \frac{1}{\varepsilon}}{2h}\right)^{n-\alpha+\frac{1}{2}} J_{n-\alpha+\frac{1}{2}}\left(h \ln \frac{1}{\varepsilon}\right) (h^2 + L_0^2)^{n+1} [f](\varepsilon z) d\varepsilon, \end{aligned}$$

i.e., is an integral-differential operator.

3.7. We let

$$\Delta(p) = \sqrt{h + p e^{\frac{k}{h+p}}}; \quad k, h > 0.$$

In the table of the Laplace transformations we choose the formula

$$e^{-ht} \frac{1}{\sqrt{\pi t}} \cos 2\sqrt{kt} \rightarrow \frac{1}{\sqrt{h + x e^{\frac{k}{h+x}}}.$$

Due to Theorem 2.1 and formula (2.1) we find the expression for the operator L_ω :

$$L_\omega[f] = \frac{1}{\sqrt{\pi}} \int_0^1 \frac{\varepsilon^{h-1}}{\sqrt{\ln \frac{1}{\varepsilon}}} \cos 2\sqrt{k \ln \frac{1}{\varepsilon}} f(\varepsilon z) d\varepsilon.$$

Taking into consideration the formula

$$e^{-ht} \frac{1}{\sqrt{\pi t}} ch2\sqrt{kt} \rightarrow \frac{1}{\sqrt{h+x}} e^{\frac{k}{h+x}},$$

we find the inverse operator L_{ω}^{-1} :

$$L_{\omega}^{-1} [f] = \frac{1}{\sqrt{\pi}} (h + L_0) \int_0^1 \frac{\varepsilon^{h-1}}{\sqrt{\ln \frac{1}{\varepsilon}}} ch2\sqrt{k \ln \frac{1}{\varepsilon}} \cdot f(\varepsilon z) d\varepsilon.$$

4. CONCLUSION

In the work we presented the technique of applying integral Laplace transform in the theory of the generalized Riemann-Liouville operators. We found the closed formulae for the inverse of the generalized Riemann-Liouville operators. The main result of the work is the extension of the scalar sequence

$$\Delta(k) = k \int_0^1 r^{k-1} \omega(x) dx < \infty, k = 1, 2, \dots$$

in the half-plane $p : \operatorname{Re} p = \sigma > \sigma_0 \geq 0$. The problem on inversion of the generalized Riemann-Liouville operator is solved in the case when the function $\Delta(p)$ is the Laplace image, while the function $\Delta^{-1}(p)$ has a power growth, i.e.,

$$\frac{1}{\Delta(p)} = p^{\alpha} L(p), \quad \alpha > 0,$$

where $L(p)$ is a slowly varying function [14].

In particular, in the work we presented the theory of fractional powers of the Hadamard operator; in our notations, this is the operator L_0 . We considered the theory of the fractional powers of the generalized Hadamard operators such as the operator $h + L_0$, $h > 0$, and the operator $h^2 + L_0^2$, $h > 0$.

BIBLIOGRAPHY

1. I.I. Bavrin. *The operator method in complex analysis*. Prometeĭ, Moscow (1990). (in Russian).
2. I.I. Bavrin, V.L. Matrosov, O.E. Yaremko. *Integral transforms and representations of functions in real and complex domains*. Prometeĭ, Moscow (2000). (in Russian).
3. A.S. Berdyshev, B.Kh. Turmetov, B.J. Kadirkulov. *Some properties and applications of the integrodifferential operators of Hadamard-Marchaud type in the class of harmonic functions // Sibir. Matem. Zhurn.* **53**:4, 752–764 (2012). [Siber. Math. J. **53**:4, 600–610 (2012).]
4. A. Erdélyi, W. Magnus, F. Oberhettinger, F.G. Tricomi. *Higher transcendental functions. V. I*. McGraw-Hill Book Co., New York (1953).
5. M.M. Džrbašjan. *A generalized Riemann-Liouville operator and some of its applications // Izv. AN SSSR. Ser. Matem.* **32**:5, 1075–1111 (1968). [Math. USSR. Izv. **2**:5, 1027–1063 (1968).]
6. M.M. Dzhrbashyan, A.B. Nersesjan. *Fractional derivatives and Cauchy problem for fractional order differential equations // Izv. AN Armyan. SSR. Ser. Matem.* **3**:1, 3–29 (1968). (in Russian).
7. A.M. Nakhushev. *Fractional calculus and its applications*. Fizmatlit, Moscow (2003). (in Russian.)
8. A.M. Nakhushev. *Equations of mathematical biology*. Vysshaya Shkola, Moscow (1995). (in Russian).
9. A.A. Potapov. *Brief historical essay on arising and formation of the theory of fractional integration-differentiation // Nelinejnyj mir.* **1**:1-2, 69–81 (2003). (in Russian).
10. St.G. Samko, A.A. Kilbas, O.I. Marichev. *Fractional integrals and derivatives: theory and applications*. Nauka i tekhnika, Minsk (1987). [Gordon and Breach, New York (1993).]
11. V.V. Uchaikin. *Method of fractional derivatives*. Artishok, Ul'yanovsk (2008). (in Russian.)
12. E. Hille, R.S. Einar. *Functional analysis and semigroups*. Colloquium Publications. **31**. Amer. Math. Soc. Providence, RI. (1957).

13. V.A. Churikov. *Fractional analysis on the base of the Hadamard operator* // Izvestia Tomsk. Politekh. Univ. **312**:2, 16–20 (2008). (in Russian).
14. W. Feller. *An introduction to probability theory and its applications. V. II.* Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, New York. (1971).
15. A.A. Kilbas, H.M. Srivastava., J.J. Trujillo. *Theory and applications of fractional differential equations.* North-Holland Mathematics Studies **204**. Elsevier, Amsterdam (2006).
16. A.N. Kochubei. *Distributed order derivatives and relaxation patterns* // J. Phys. A: Math. Theor. **42**:31, id 315203 (2009).
17. A.N. Kochubei. *Distributed order calculus and equations of ultraslow diffusion* // J. Math. Anal. Appl. **340**:1, 251–282 (2008).
18. A.N. Kochubei. *General fractional calculus, evolution equations, and renewal processes* // Integ. Equat. Oper. Theor. **71**:4, 583–600 (2011).

Ivan Ivanovich Bavrin,
Moscow State Pedagogical University,
Malaya Pirogovskaya str. 1,
119435, Moscow, Russia
E-mail: ivbavrin@yandex.ru

Oleg Emanuilovich Yaremko,
Penza State University,
Krasnaya str. 40,
440038, Penza, Russia
E-mail: yaremki@mail.ru