

doi:10.13108/2021-13-3-174

PERTURBATION THEORY METHODS IN PROBLEM OF PARAMETRIC RESONANCE FOR LINEAR PERIODIC HAMILTONIAN SYSTEMS

M.G. YUMAGULOV, L.S. IBRAGIMOVA, A.S. BELOVA

Abstract. We consider the problem on parametric resonance for linear periodic Hamiltonian systems depending on a small parameter. We propose new formulae based on the methods of the perturbation theory for linear operators in the problem on approximate construction of multipliers for linear non-autonomous periodic Hamiltonian systems. We focus on obtaining the formulae for the first correctors of perturbations of multiple definite and indefinite multipliers. The proposed formulae lead to new Lyapunov stability criteria for linear periodic Hamiltonian systems in critical cases. We consider applications to the problem on parametric resonance in main resonances. The obtained results are formulated in terms of the original equations and lead us to effective formulae and algorithms. The effectiveness of the proposed formulae is demonstrated by solving the problem of plotting the boundaries of the stability regions of triangular libration points of a planar bounded elliptic three-body problem.

Keywords: Hamiltonian system, stability, multiplier, small parameter, parametric resonance, perturbation theory, three-body problem, libration point.

Mathematics Subject Classification: 37J25, 37J40, 37N05

1. INTRODUCTION

1.1. Main equations. We consider a linear periodic Hamiltonian system (LPHS) depending on small scalar or vector parameter ε :

$$\frac{dx}{dt} = JA(t, \varepsilon)x, \quad x \in \mathbb{R}^{2N}, \quad (1.1)$$

in which $A(t, \varepsilon)$ is a real symmetric T -periodic in t matrix, that is, $A(t + T, \varepsilon) \equiv A(t, \varepsilon)$, while the matrix J is defined by the identity:

$$J = \begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix}; \quad (1.2)$$

here I is the unit $N \times N$ matrix. We assume that

- the entries of the matrix $A(t, \varepsilon)$ are continuous in t and \mathbb{C}^k -smooth in ε , where $k \geq 1$;
- the identity holds:

$$A(t, 0) \equiv A_0, \quad (1.3)$$

where A_0 is a constant symmetric matrix.

M.G. YUMAGULOV, L.S. IBRAGIMOVA, A.S. BELOVA, PERTURBATION THEORY METHODS IN PROBLEM OF PARAMETRIC RESONANCE FOR LINEAR PERIODIC HAMILTONIAN SYSTEMS.

© YUMAGULOV M.G., IBRAGIMOVA L.S., BELOVA A.S. 2021.

The research of the third author is made in the framework of State Task of the Ministry of Science and Higher Education of Russian Federation (code of scientific theme FZWU-2020-0027).

Submitted February 18, 2021.

Condition (1.3) that as $\varepsilon = 0$, LPHS (1.1) is a linear autonomous Hamiltonian system (LAHS):

$$\frac{dx}{dt} = JA_0x, \quad x \in \mathbb{R}^{2N}. \tag{1.4}$$

We call system (1.4) unperturbed, while (1.1) is called perturbed system.

In the paper we discuss some issues on constructing the formulae for the first corrector while perturbing multiple definite and indefinite multipliers of system (1.4). We consider their applications to studying the problem on a parametric resonance in system (1.1). A more detailed formulation will be given in what follows.

1.2. Auxiliary facts. We first briefly some auxiliary facts from the general theory of linear Hamiltonian systems, see, for instance, [1]–[4].

We consider a linear periodic Hamiltonian system:

$$\frac{dx}{dt} = JA(t)x, \quad x \in \mathbb{R}^{2N}; \tag{1.5}$$

here $A(t)$ is a real symmetric matrix, the entries of which are continuous and T -periodic in t functions. In what follows, the matrix $JA(t)$ involved in system (1.5) will be called, for simplicity, *Hamiltonian*; in literature, other terms are also used, for instance, *infinitesimal symplectic* matrix.

The multipliers of system (1.5) are the eigenvalues of its monodromy matrix V , that is, of the matrix $V = X(T)$, where $X(t)$ is a fundamental matrix of solutions (FMS) of system (1.5). As a FMS of system (1.5), we consider the solution to the matrix Cauchy problem: $X' = JA(t)X$, $X(0) = I$; here I is the identity matrix of dimension $2N$.

The following facts hold:

- Let LPHS (1.5) possesses a multiplier μ_0 . Then $\mu_0 \neq 0$ and the numbers $\frac{1}{\mu_0}$, $\bar{\mu}_0$, $\frac{1}{\bar{\mu}_0}$ are also the multipliers of LPHS (1.5) and they are of the same algebraic and geometric multiplicity and are of the same index.
- If LPHS (1.5) has a multiplier 1 (or -1), then this multiplier has an even algebraic multiplicity.
- LPHS (1.5) is stable if and only if all its multipliers μ are located on the unit circumference, that is, $|\mu| = 1$, and are semi-simple. At that, LPHS (1.5) is not necessarily asymptotically stable.

Together with (1.5), we shall also consider a perturbed LPHS of form:

$$\frac{dx}{dt} = J\tilde{A}(t)x, \quad x \in \mathbb{R}^{2N}, \tag{1.6}$$

in which $\tilde{A}(t)$ is a small perturbation of the matrix $A(t)$ in the class of symmetric, continuous and T -periodic matrices.

Let all multipliers μ of system (1.5) be located on the unit circle, that is, $|\mu| = 1$, and be simple. Let μ_0 be one of these multipliers. Then for small perturbations system (1.6) also has a simple multiplier $\tilde{\mu}$ close to μ_0 and $|\tilde{\mu}| = 1$. The stability property of system (1.5) in this case do not change: it remains stable.

Now let system (1.5) has a multiple multiplier μ_0 such that $|\mu_0| = 1$ and other multipliers are still simple and are located on the unit circumference. For simplicity, let this multiplicity be 2. Then while passing from (1.5) to a perturbed system (1.6), the multiplier μ_0 usually splits into two simple multipliers μ_1 and μ_2 . At that, two cases are possible:

- a) $\mu_2 = 1/\overline{\mu_1} : |\mu_1| < 1 < |\mu_2|;$
 b) $|\mu_1| = |\mu_2| = 1.$

In Case a), perturbed system (1.6) becomes unstable, while in Case b) it is stable.

Rigorously speaking, the question on which of Cases a) or b) is chosen by system (1.5) under the splitting of a multiple multiplier μ_0 depends not only on the properties of perturbed system (1.6) but also on unperturbed system (1.5). Namely, multiple multipliers are divided into two types: *definite* and *indefinite*, see, for instance, [2]. If a multiple multiplier μ_0 is definite, then for each sufficiently small linear periodic Hamiltonian perturbation of system (1.5) the multiplier μ_0 can split only via Case b), that is, the multiplier μ_1 and μ_2 stay on the unit circumference. If the multiple multiplier μ_0 is indefinite, then there exist small perturbations which move these multipliers away from the unit circumference.

The notion of the definiteness of the multipliers is related with another important notion of the theory of LPHS. We say, see, for instance, [2], [5], that system (1.5) is *strongly (parametrically) stable* if it and all its small linear periodic Hamiltonian perturbations are stable in the Lyapunov sense. In other words, system (1.5) is strongly (parametrically) stable if this system and close to it perturbed systems (1.6) are stable.

The following statement is of an important value, see, for instance, [2], [5].

Theorem 1.1 (Krein-Gelfand-Lidskii). *System (1.5) is strongly stable if and only if*

- 1) *all its multipliers are semi-simple and their absolute values are equal to one;*
- 2) *the numbers ± 1 are not its multipliers;*
- 3) *all its multiple multipliers are definite.*

2. FORMULATION OF PROBLEM

We return back to main systems (1.1) and (1.4).

For unperturbed autonomous system (1.4), the monodromy matrix V in the T -periodic problem reads as $V = e^{TJA_0}$. The multipliers μ of system (1.4) are related with the eigenvalues λ of the matrix JA_0 by the identity $\mu = e^{T\lambda}$. By the aforementioned properties of linear Hamiltonian system and in accordance with the perturbation theory of linear operators, see, for instance, [19]), the following holds: if the matrix JA_0 possesses at least one eigenvalue with a non-zero real part, then perturbed LPHS (1.1) is unstable for all small $|\varepsilon|$.

Let all eigenvalues of the matrix JA_0 be pure imaginary, namely, these are numbers

$$\pm i\omega_1, \quad \pm i\omega_2, \quad \dots, \quad \pm i\omega_N, \quad (2.1)$$

where $\omega_j \geq 0$. If the algebraic multiplicity of some eigenvalue $i\omega_m$ is k , then in list (2.1) the number $i\omega_m$ appears exactly k times. In this case, as $\varepsilon = 0$, the absolute values of all multipliers of LPHS (1.1) are equal to one. As it has been mentioned above, the situations, when some of these multipliers are multiple, are of a special interest.

Multiple multipliers of system (1.4) arise under one of the following condition, see, for instance [1]):

S1) among number (2.1), there exists at least one $i\omega_{m_0}$ such that

$$\omega_{m_0} = \frac{\pi k_0}{T} \quad \text{for some integer non-negative } k_0; \quad (2.2)$$

S2) among numbers (2.1) there exists at least one pair $i\omega_{m_0}$ and $i\omega_{l_0}$ ($m_0 \neq l_0$) such that

$$\omega_{m_0} - \omega_{l_0} = \frac{2\pi k_0}{T} \quad \text{for some integer } k_0. \quad (2.3)$$

Remark 2.1. Identity (2.2) means that the corresponding multiplier of system (1.4) is equal to 1 if k_0 is even or -1 if k_0 is odd and this multiplier is of an even multiplicity. By Krein-Gelfand-Lidskii theorem this implies that under identity (2.2) unperturbed system (1.4) does not possess the property of strong stability.

Remark 2.2. Identity (2.3) means that the corresponding multiplier of system (1.4) is multiple and it equals to

$$\mu_0 = e^{T\omega_{m_0}i} = e^{T\omega_{l_0}i}. \quad (2.4)$$

If at the same time the numbers ω_{m_0} and ω_{l_0} do not satisfy relations of form (2.2), that is, $\omega_{m_0}, \omega_{l_0} \neq \pi k/T$ for all integer k , then multiplier (2.4) obeys the inequality $\mu_0 \neq \pm 1$.

Condition **S2** covers also the case when the matrix JA_0 has a multiple pure imaginary eigenvalue. Namely, this is the case if identity (2.3) holds with $k_0 = 0$: then $\omega_{m_0}i = \omega_{l_0}i$.

The problem on stability of system (1.1) under conditions of type **S1** or **S2** is often called *problem on parametric resonance*, see, for instance, [1], [7], and the relations are called *parametric resonances*. Relation (2.2) is called a *simple resonance*, while a relation of type (2.3) is called *combinational resonance*. In view of this we mention that relations (2.2) and (2.3) can be represented in an universal form:

$$n_1\omega_{m_0} + n_2\omega_{l_0} = \frac{2\pi k_0}{T},$$

in which n_1 and n_2 are integer numbers such that $|n_1| + |n_2| = 2$. In this case one says about a *second order resonance*.

A lot of works are devoted to studying the stability of linear Hamiltonian systems with a periodic perturbation, in particular, to the problem on parametric resonance. The most part of studies are based on renormalization methods of linear Hamiltonian systems and on a transformation of the Hamiltonian of system (1.1) by means of a canonical change of variables. In this direction, a series of important results was obtained, see, for instance, [1], [11]–[17].

Other approaches to studying the problem on parametric resonance are based on the classical perturbation theory of linear operators. It should be said that direct application of the methods of general theory is not of interest since this way, as a rule, is extremely cumbersome and therefore is not applied in practice. At the same, the main interest is related with the modifications of these methods, which as much as possible take into consideration the features of the problem associated with the Hamiltonian property of the systems. This approach was developed in works by many authors, see, for instance, [2], [5]–[8].

Nowadays the researches are continued in various directions. It is most topical to develop general approaches for investigating the problem on parametric resonance in terms of the original equations without the need of preliminary transformation, which is often laborious and cumbersome. The main difficulty is the problem of constructing formulae for the first correctors of the multipliers of a perturbed non-autonomous periodic Hamiltonian system. In this regard, we point out that the formulae known in the literature, as a rule, are aimed at studying autonomous systems, see, for instance, [5], [8], [10], [18].

In the present paper we study the problem on parametric resonance for LPHS (1.1) under Conditions **S1** and **S2**. We provide new formulae for the first correctors while approximately constructing the multipliers for system (1.1) taking into consideration their definiteness or indefiniteness. The obtained formula are used for analyzing the stability of LPHS (1.1) in the Lyapunov sense.

The problem on parametric resonance of LPHS (1.1) is studied in the following main cases corresponding to conditions **S1** and **S2**:

P_1 . The matrix JA_0 has a multiple (double) eigenvalue $\lambda = i\omega_0$, where $\omega_0 \geq 0$ and $\omega_0 \neq \pi k/T$ for natural k .

P_2 . The matrix JA_0 possesses two simple eigenvalues $\lambda_1 = i\omega_1$ and $\lambda_2 = i\omega_2$, where $\omega_1, \omega_2 > 0$, $\omega_1, \omega_2 \neq \pi k/T$ for natural k and $\omega_1 - \omega_2 = 2\pi k_0/T$ for some natural k_0 .

P_3 . The matrix JA_0 has a simple eigenvalue $\lambda = i\omega_0$, where $\omega_0 = \pi k_0/T$ for some natural k_0 .

We assume that the other eigenvalues λ of the matrix JA_0 , different from $\pm i\omega_0$ in Cases P_1 and P_3 and from $\pm i\omega_1$ and $\pm i\omega_2$ in Case P_2 , are simple and pure imaginary, namely, they are of form $\lambda = i\omega$, where $\omega \neq \pi k/T$ for integer k . At the same time, none of these pairs satisfies a resonance relation like one given in Case P_2 .

Cases P_1 and P_2 correspond to condition (2.3), while Case P_3 does to condition (2.2). However, if in Case P_1 we have $\omega_0 = 0$, then it corresponds to both conditions (2.2) and (2.3).

It is convenient to represent system (1.1) in the form

$$\frac{dx}{dt} = J[A_0 + \varepsilon S_1(t) + S_2(t, \varepsilon)]x, \quad x \in \mathbb{R}^{2N}, \quad (2.5)$$

where J is matrix (1.2), A_0 is matrix (1.3), $S_1(t)$ and $S_2(t, \varepsilon)$ are real symmetric and T -periodic in t matrices, and $S_2(t, \varepsilon)$ is smooth in ε and satisfies the relation $\|S_2(t, \varepsilon)\| = O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$ uniformly in t .

3. CASE P_1

We begin studying of problem from Case P_1 . This case is partitioned into subcases, when the eigenvalue $\lambda = i\omega_0$ is semi-simple (subcase P_1^1) or not (subcase P_1^2). We proceed to considering these subcases.

3.1. Subcase P_1^1 . Let the matrix JA_0 has a semi-simple (double) eigenvalue $i\omega_0$, where $\omega_0 \geq 0$ and $\omega_0 \neq \pi k/T$ for natural k . We denote by $V(\varepsilon)$ the monodromy matrix of perturbed system (2.5). Then $V_0 = e^{JA_0 T}$ is the monodromy matrix of unperturbed matrix (1.4). In the considered subcase the matrix V_0 has a semi-simple eigenvalue $\mu_0 = e^{i\omega_0 T}$ of multiplicity 2. We note that since $\omega_0 \neq \pi k/T$ for natural k , then for $\omega_0 > 0$ we have the inequality $\mu_0 \neq \pm 1$. We also note that as $\omega_0 > 0$ the multiplier μ_0 of system (1.4) can be both definite and indefinite, while as $\omega_0 = 0$, it is indefinite.

Namely, as $\omega_0 > 0$, two mutually exclusive cases are possible, see, for instance, [2], [5]:

- 1⁰ for each eigenvector e associated with the eigenvalue $i\omega_0$ of the matrix JA_0 the relation holds: $(Je, e) \neq 0$;
- 2⁰ there exists an eigenvector e associated with the eigenvalue $i\omega_0$ of the matrix JA_0 , for which $(Je, e) = 0$.

In the first case the multiplier $\mu_0 = e^{i\omega_0 T}$ of system (1.4) is definite, while in the second case it is indefinite.

Since the matrix JA_0 has a semi-simple (double) eigenvalue $i\omega_0$, then there exist non-zero linearly independent vectors $e, g \in \mathbb{C}^{2N}$ such that

$$JA_0 e = i\omega_0 e, \quad JA_0 g = i\omega_0 g. \quad (3.1)$$

The vectors e, g are the eigenvectors also for the monodromy matrix $V_0 = e^{JA_0 T}$ and are associated with a semi-simple eigenvalue $\mu_0 = e^{i\omega_0 T}$ of multiplicity 2.

The aforementioned properties of the multipliers μ_0 related with the definiteness and indefiniteness imply the following statement.

Lemma 3.1. *Let μ_0 be a definite multiplier of system (1.4). Then the eigenvectors e, g of the matrix JA_0 can be normalized only by one of the following identities:*

$$(iJe, e) = (iJg, g) = 1 \quad (3.2)$$

or

$$(iJe, e) = (iJg, g) = -1. \quad (3.3)$$

If μ_0 is an indefinite multiplier, then the vectors e, g can be normalized by the identities

$$(iJe, e) = -1, \quad (iJg, g) = 1. \tag{3.4}$$

Here the vectors e, g can be chosen satisfying the identity:

$$(e, Jg) = 0. \tag{3.5}$$

According to the perturbation theory of linear operators, see, for instance, [19]), for small $|\varepsilon|$ the matrix $V(\varepsilon)$ has a pair of eigenvalues $\mu_1(\varepsilon)$ and $\mu_2(\varepsilon)$ such that the functions $\mu_1(\varepsilon)$ and $\mu_2(\varepsilon)$ are continuously differentiable, and $\mu_1(0) = \mu_2(0) = \mu_0$. Moreover, they can be represented in the form

$$\mu_1(\varepsilon) = \mu_0 + \mu_1^{(1)}\varepsilon + O(\varepsilon^{3/2}), \quad \mu_2(\varepsilon) = \mu_0 + \mu_1^{(2)}\varepsilon + O(\varepsilon^{3/2}). \tag{3.6}$$

Let us provide the scheme of constructing the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in formulae (3.6).

Let the vectors e, g be normalized in accordance with Lemma 3.1. We define a constant matrix

$$S_0 = \int_0^T S_1(t) dt, \tag{3.7}$$

where $S_1(t)$ is a matrix involved in system (2.5). We let

$$a = (S_0e, e), \quad b = (S_0g, g), \quad c = (S_0g, e). \tag{3.8}$$

We note that the numbers a and b are real, while the number c is in general complex.

3.1.1. Perturbation of definite multiplier. We first consider the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in formulae (3.6) under the assumption that $\mu_0 = e^{i\omega_0 T}$ is a definite multiplier of system (1.4). We note that then $\mu_0 \neq \pm 1$ and therefore, in formula $\mu_0 = e^{i\omega_0 T}$ we necessarily have $\omega_0 \neq \pi k/T$ for integer k . In particular, here we should assume that $\omega_0 \neq 0$.

Theorem 3.1. *Let one of normalizations (3.2) and (3.3) hold and identity (3.5) be satisfied. Then the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in expansions (3.6) are the eigenvalues of the matrices $D = -i\mu_0 D_0$ and $D = i\mu_0 D_0$ respectively, where*

$$D_0 = \begin{bmatrix} a & c \\ \bar{c} & b \end{bmatrix}. \tag{3.9}$$

The proof of this and other main statements of the paper are given in Section 7.

We note that the eigenvalues λ_1 and λ_2 of matrix (3.9) are real numbers, namely, they are roots of the quadratic equation

$$\lambda^2 - (a + b)\lambda + ab - c\bar{c} = 0. \tag{3.10}$$

We provide some corollaries from Theorem 3.1.

Corollary 3.1. *Let μ_0 be a definite multiplier of system (1.4). Let normalization (3.2) or normalization (3.3) hold. Then the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in expansions (3.6) read as*

$$\mu_1^{(1)} = -i\mu_0\lambda_1, \quad \mu_1^{(2)} = -i\mu_0\lambda_2 \quad (\mu_1^{(1)} = i\mu_0\lambda_1, \quad \mu_1^{(2)} = i\mu_0\lambda_2), \tag{3.11}$$

where λ_1 and λ_2 are the roots of quadratic equation (3.10).

Corollary 3.2. *Let μ_0 be a definite multiplier of system (1.4) and normalization (3.2) or (3.3) hold. Then for $\omega_0 > 0$ unperturbed system (1.4) is strongly stable. While passing to perturbed system (1.1), the definite multiplier μ_0 splits in accordance with formulae (3.6) and (3.11) staying on the unit circumference: $|\mu_1(\varepsilon)| = |\mu_2(\varepsilon)| = 1$. For small $|\varepsilon|$ system (1.1) is stable.*

3.1.2. *Perturbation of indefinite multiplier.* Now let $\mu_0 = e^{i\omega_0 T}$ be an indefinite multiplier of system (1.4).

Theorem 3.2. *Let normalization (3.4) and identity (3.5) hold. Then the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in expansions (3.6) are the eigenvalues of the matrix $D = i\mu_0 D_1$, where*

$$D_1 = \begin{bmatrix} a & c \\ -\bar{c} & -b \end{bmatrix}. \quad (3.12)$$

The eigenvalues λ_1 and λ_2 of the matrix (3.12) are the roots of the quadratic equation

$$\lambda^2 + (a - b)\lambda - ab + c\bar{c} = 0 \quad (3.13)$$

and therefore, they can be both real and complex. We denote by $\Delta = (a + b)^2 - 4c\bar{c}$ the discriminant of equation (3.13).

We provide corollaries from Theorem 3.2.

Corollary 3.3. *Let μ_0 be an indefinite multiplier of system (1.4) and normalization (3.4) hold. Then the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in expansions (3.6) read as*

$$\mu_1^{(1)} = -i\mu_0\lambda_1, \quad \mu_1^{(2)} = -i\mu_0\lambda_2, \quad (3.14)$$

where λ_1 and λ_2 are the roots of quadratic equation (3.13). For small $|\varepsilon|$, the indefinite multiplier μ_0 of system (1.4) splits according to formulae (3.6) and (3.14).

Corollary 3.4. *Let $\Delta > 0$. Then for a given perturbation $S_1(t)$ of system (2.5) and all small $|\varepsilon|$ the indefinite multiplier μ_0 of system (1.4) splits but stays on the unit circle: $|\mu_1(\varepsilon)| = |\mu_2(\varepsilon)| = 1$. For small $|\varepsilon|$ system (2.5) is stable.*

Corollary 3.5. *Let $\Delta < 0$. Then for a given perturbation $S_1(t)$ of system (2.5) and all small non-zero $|\varepsilon|$ the indefinite multiplier μ_0 of system (1.4) splits leaving the unit circle: $|\mu_1(\varepsilon)| < 1$ and $|\mu_2(\varepsilon)| > 1$. For small non-zero $|\varepsilon|$ system (2.5) is unstable.*

3.2. Subcase P_1^2 . Let the matrix JA_0 possesses a non-semi-simple (double) eigenvalue $i\omega_0$, where $\omega_0 \geq 0$ and $\omega_0 \neq \pi k/T$ for natural k . Then the monodromy matrix V_0 of unperturbed system (1.4) has a non-semi-simple eigenvalue $\mu_0 = e^{\omega_0 T i}$ of multiplicity 2. According to the perturbation theory of linear operator, for small $|\varepsilon|$ the monodromy matrix $V(\varepsilon)$ of perturbed system (2.5) has a pair of eigenvalues $\mu_1(\varepsilon)$ and $\mu_2(\varepsilon)$ such that the functions $\mu_1(\varepsilon)$ and $\mu_2(\varepsilon)$ are continuous and $\mu_1(0) = \mu_2(0) = \mu_0$. Moreover, they can be represented by Puiseux series:

$$\mu_1(\varepsilon) = \mu_0 + \mu_1^{(1)}\varepsilon^{1/2} + O(\varepsilon), \quad \mu_2(\varepsilon) = \mu_0 + \mu_2^{(1)}\varepsilon^{1/2} + O(\varepsilon). \quad (3.15)$$

We proceed to calculating the coefficients $\mu_1^{(j)}$ in formulae (3.15). We observe that in the considered case there exists a pair of non-zero linearly independent vectors $e, g \in \mathbb{C}^{2N}$ such that the identities hold:

$$JA_0 e = i\omega_0 e, \quad JA_0 g = i\omega_0 g + e. \quad (3.16)$$

We also mention the identities $V_0 e = \mu_0 e$ and $V_0 g = \mu_0(g + Te)$.

It is easy to prove the following statement.

Lemma 3.2. *The relations $(e, Je) = 0$, $(e, Jg) \neq 0$ hold and the number (e, Jg) is real. The vector g can be chosen by the identity*

$$(g, Jg) = 0. \quad (3.17)$$

In what follows we suppose that identity (3.17) holds. By Lemma 3.2, the number $\nu = \frac{1}{(e, Jg)}$ is well-defined.

Theorem 3.3. *The coefficients $\mu_1^{(1)}$ and $\mu_2^{(1)}$ in expansions (3.15) read as*

$$\mu_1^{(1)} = \mu_0 \sqrt{-T\nu(S_0 e, e)}, \quad \mu_2^{(1)} = -\mu_1^{(1)}, \quad (3.18)$$

where S_0 is matrix (3.7).

We note that the radicand in formula (3.18) is real.

We provide some corollaries from Theorem 3.3.

Corollary 3.6. *In Subcase P_1^2 unperturbed system (1.4) is not strongly stable and respectively, its multiplier $\mu_0 = e^{i\omega_0 T}$ is indefinite. For small $|\varepsilon|$ the multiplier μ_0 splits in accordance with formulae (3.15) and (3.18).*

Corollary 3.7. *Let $\varepsilon\nu(S_0 e, e) > 0$. Then for a given perturbation $S_1(t)$ of system (2.5), for small $|\varepsilon|$, the multiplier μ_0 of system (1.4) stays on the unit circumference: $|\mu_1(\varepsilon)| = |\mu_2(\varepsilon)| = 1$. In this case system (2.5) is stable.*

Corollary 3.8. *Let $\varepsilon\nu(S_0 e, e) < 0$. Then for a given perturbation $S_1(t)$ of system (2.5) for corresponding small $|\varepsilon|$, the multiplier μ_0 of system (1.4) leaves the unit circumference $|\mu_1(\varepsilon)| < 1$ and $|\mu_2(\varepsilon)| > 1$. In this case system (2.5) is unstable.*

3.3. Case of zero eigenvalue. An important particular version of Case P_1 is a situation, when the matrix JA_0 has a zero eigenvalue $\lambda = 0$ of multiplicity 2. Here there are some features.

We first mention that in this situation in formulae (3.6) and (3.15) we have $\mu_0 = 1$. By Krein-Gelfand-Lidskii this implies that then system (1.4) does not possess the property of strong stability.

Second, the eigenvalue $\lambda = 0$ can be semi-simple or not. This is this situation will be studied in accordance with the formulae and conclusions of Subsections 3.1.2 and 3.2.

4. CASE P_2

Here we consider problem on parametric resonance (2.5) in Case P_2 , that is, as JA_0 two simple eigenvalues $\lambda_1 = i\omega_1$ and $\lambda_2 = i\omega_2$, where $\omega_1, \omega_2 > 0$, $\omega_1, \omega_2 \neq \pi k/T$ for natural k and $\omega_1 - \omega_2 = 2\pi k_0/T$ for some natural k_0 . Let, as above, $V(\varepsilon)$ is the monodromy matrix of perturbed system (2.5). Then the monodromy matrix $V_0 = e^{JA_0 T}$ of unperturbed system (1.4) has a semi-simple eigenvalue μ_0 of multiplicity 2, namely, $\mu_0 = e^{T\omega_1 i} = e^{T\omega_2 i}$. At that, since $\omega_1, \omega_2 \neq \pi k/T$ for natural k , we have $\mu_0 \neq \pm 1$.

For small $|\varepsilon|$ the matrix $V(\varepsilon)$ has two eigenvalues $\mu_1(\varepsilon)$ and $\mu_2(\varepsilon)$ such that the functions $\mu_1(\varepsilon)$ and $\mu_2(\varepsilon)$ are continuously differentiable and can be represented in form (3.6).

Since in considered Case P_2 the multiplier μ_0 of system (1.4) is semi-simple, the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in formulae (3.6) can be constructed by the same scheme as for Cubcase P_1^1 , see Subsection 3.1. Here we mention just some features related to Case P_2 .

First, instead of identities (3.1), which hold in subcase P_1^1 , in Case P_2 there exists a pair of linearly independent vectors $e, g \in \mathbb{C}^{2N}$ such that the identities

$$JA_0 e = i\omega_1 e, \quad JA_0 g = i\omega_2 g$$

are true. These vectors are also eigenvectors of the monodromy matrix $V_0 = e^{JA_0 T}$ associated with the semi-simple double eigenvalue μ_0 .

Second, although here a complete analogue of Lemma 3.1 holds true, the proof of this analogue has its own features. In particular, there is no need to ensure identity (3.5) since now it is always true.

Finally, third, instead of numbers (3.8) in matrices (3.9) and (3.12) in Case P_2 we should employ the numbers

$$a = (S_0e, e), \quad b = (S_0g, g), \quad c = \int_0^T e^{-2\pi i k_0 t/T} (S_1(t)g, e) dt; \tag{4.1}$$

here S_0 is matrix (3.7).

In view of the mentioned features in the considered Case P_2 , complete analogues of Theorems 3.1 and 3.2 hold as well as of their corollaries.

We note that as $k_0 = 0$, the formulae for Case P_2 coincide with the corresponding formulae of the above considered Subcase P_1^1 . This is natural since the identity $k_0 = 0$ means that $\omega_1 = \omega_2$, that is, the matrix JA_0 has a multiple semi-simple eigenvalue $\lambda = i\omega_0$; here $\omega_0 = \omega_1 = \omega_2$.

5. CASE P_3

We proceed to Case P_3 . Let the matrix JA_0 has a simple eigenvalue $i\omega_0$, where $\omega_0 = \pi k_0/T$ for some natural k_0 . Then the monodromy matrix $V_0 = e^{JA_0T}$ of unperturbed system (1.4) has a semi-simple eigenvalue μ_0 of multiplicity 2, where $\mu_0 = 1$ if k_0 is even and $\mu_0 = -1$ if k_0 is odd. It follows from Krein-Gelfand-Lidskii theorem that in the considered case system (1.4) does not possess the property of strong stability.

For small $|\varepsilon|$, the monodromy matrix $V(\varepsilon)$ of perturbed system (2.5) possesses a pair of eigenvalues $\mu_1(\varepsilon)$ such that $\mu_2(\varepsilon) \mu_1(0) = \mu_2(0) = \mu_0$. The functions $\mu_1(\varepsilon)$ and $\mu_2(\varepsilon)$ are continuously differentiable and are represented in form (3.6).

Let us provide a statement about calculating the coefficients $\mu_1^{(j)}$ in formulae (3.6). We mention that in the considered case there exists a non-zero vector $e+ig \in \mathbb{C}^{2N}$, where $e, g \in \mathbb{R}^{2N}$ such that

$$JA_0(e + ig) = i\omega_0(e + ig). \tag{5.1}$$

The vectors $e, g \in \mathbb{R}^{2N}$ turn out to be eigenvectors also for the monodromy matrix $V_0 = e^{JA_0T}$ associated with a semi-simple eigenvalue μ_0 of multiplicity 2.

Lemma 5.1. *The inequality $(e, Jg) \neq 0$ holds.*

We let

$$\nu = \frac{1}{(e, Jg)}. \tag{5.2}$$

The number (e, Jg) , and therefore, the number ν are real. We define a matrix:

$$B = \nu\mu_0 \begin{bmatrix} a & b_1 \\ b_2 & -a \end{bmatrix}, \tag{5.3}$$

in which the numbers a, b_1 and b_2 are given by the identities:

$$a = \int_0^T \{ \cos(2\omega_0 t) (S_1(t)e, g) - \frac{1}{2} \sin(2\omega_0 t) [(S_1(t)g, g) - (S_1(t)e, e)] \} dt, \tag{5.4}$$

$$b_1 = \int_0^T [\cos^2(\omega_0 t) (S_1(t)g, g) + \sin^2(\omega_0 t) (S_1(t)e, e) + \sin(2\omega_0 t) (S_1(t)e, g)] dt, \tag{5.5}$$

$$b_2 = b_1 - [(S_0e, e) + (S_0g, g)]; \tag{5.6}$$

here S_0 is matrix (3.7).

Theorem 5.1. *The coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in formulae (3.6) are the eigenvalues of matrix (5.3).*

We let

$$\Delta = a^2 + b_1 b_2. \tag{5.7}$$

The eigenvalues λ_1 and λ_2 of matrix (5.3) are numbers $\lambda_{1,2} = \pm \nu \mu_0 \sqrt{\Delta}$, which can be both real and pure imaginary. Therefore, the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in formulae (3.6) are the numbers

$$\mu_1^{(1)} = \nu \mu_0 \sqrt{\Delta}, \quad \mu_1^{(2)} = -\mu_1^{(1)}. \tag{5.8}$$

We provide some corollaries from Theorem 5.1.

Corollary 5.1. *In Case P_3 the multiplier μ_0 of system (1.4) is equal to $\mu_0 = 1$ or $\mu_0 = -1$ and is semi-simple of multiplicity 2. This multiplier is indefinite and unperturbed system (1.4) is not strongly stable. For small $|\varepsilon|$ the multiplier μ_0 splits in accordance with formulae (3.6) and (5.8).*

Corollary 5.2. *Let $\Delta < 0$. Then for a given perturbation $S_1(t)$ of system (2.5) for small $|\varepsilon|$ the multiplier μ_0 of system (1.4) stays on the unit circumference: $|\mu_1(\varepsilon)| = |\mu_2(\varepsilon)| = 1$. In this case system (2.5) remains stable.*

Corollary 5.3. *Let $\Delta > 0$. Then for a given perturbation $S_1(t)$ of system (2.5) for small non-zero $|\varepsilon|$ the multiplier μ_0 of system (1.4) leaves the unit circumference: $|\mu_1(\varepsilon)| < 1$ and $|\mu_2(\varepsilon)| > 1$. In this case for small non-zero $|\varepsilon|$ system (2.5) is unstable.*

6. APPENDIX: STABILITY OF LIBRATION POINTS IN THREE BODIES PROBLEM

6.1. Formulation of problem. As an application we consider the stability of triangle libration points in a planar bounded elliptic three-bodies problem (PBETBP), see, for instance, [5], [11]. This problem in the linear setting gives rise to a system of differential equations:

$$\frac{dx}{dt} = JA(t, \varepsilon, \mu)x, \quad x \in \mathbb{R}^4, \tag{6.1}$$

in which J is matrix (1.2) of order 4×4 , $A(t, \varepsilon, \mu)$ is a symmetric matrix:

$$A(t, \varepsilon, \mu) = \begin{bmatrix} 1 - \frac{3}{4}\rho(t, \varepsilon) & \frac{3\sqrt{3}}{4}(2\mu - 1)\rho(t, \varepsilon) & 0 & -1 \\ \frac{3\sqrt{3}}{4}(2\mu - 1)\rho(t, \varepsilon) & 1 - \frac{9}{4}\rho(t, \varepsilon) & 1 & 0 \\ 0 & 1 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{bmatrix}.$$

Here $\rho(t, \varepsilon) = (1 + \varepsilon \cos t)^{-1}$, ε is the eccentricity of Kepler orbit ($0 \leq \varepsilon < 1$), μ is a mass parameter ($0 < \mu < 1$). System (6.1) is a linear periodic (with a period $T = 2\pi$) Hamiltonian system.

Numerous studies are devoted to the stability of triangle libration points in PBETBP. One of the most interesting issues is that on constructing the stability domains for system (6.1) in the plane of parameters (μ, ε) . The main known results are presented in monograph [11]. The researches in this directions are still active, see, for instance, [22]–[25].

On Figure 1, we draw the stability and instability domains for system (6.1) for small values of μ .

The crosshatched region corresponds to the stability. The boundary of the domain is formed by three continuous curves Γ_1 , Γ_2 and Γ_3 . These curves approach the axis μ at the points

$$\mu_0 = \frac{1}{2} - \frac{\sqrt{2}}{3} = 0,028595\dots, \quad \mu^* = \frac{1}{2} - \frac{\sqrt{69}}{18} = 0,038520\dots \tag{6.2}$$

In this section, as an illustration of the above results, we discuss the construction of tangentials to the curves Γ_1 , Γ_2 and Γ_3 at the points $(\mu_0, 0)$ and $(\mu^*, 0)$, respectively.




FIGURE 1. Stability domain of triangle libration points

6.2. Auxiliary constructions. As $\varepsilon = 0$, system (6.1) is autonomous:

$$\frac{dx}{dt} = JA_0(\mu)x, \quad x \in \mathbb{R}^4, \tag{6.3}$$

where $JA_0(\mu)$ is a Hamiltonian matrix:

$$JA_0(\mu) = \begin{bmatrix} 0 & 1 & 1 & 0 \\ -1 & 0 & 0 & 1 \\ -\frac{1}{4} & \frac{3\sqrt{3}}{4}(1 - 2\mu) & 0 & 1 \\ \frac{3\sqrt{3}}{4}(1 - 2\mu) & \frac{5}{4} & -1 & 0 \end{bmatrix}. \tag{6.4}$$

An analysis of the characteristic equation of matrix (6.4) shows that

- 1° the matrix $JA_0(\mu_0)$ has a pair of pure imaginary eigenvalues of form $\pm i\omega_0$, where $\omega_0 = 1/2$. Two other its eigenvalues are the numbers $\pm i\omega_1$, where $\omega_1 = \sqrt{3}/2$. Thus, as $\mu = \mu_0$, condition of type (2.2) holds, or, what is the same, the conditions of Case P_3 , see Subsection 5. As $\mu = \mu_0$, system (6.3) has a multiplier $\eta_0 = e^{i\omega_0 2\pi} = -1$ being semi-simple of multiplicity two and indefinite.
- 2° the matrix $JA_0(\mu^*)$ has a pair of non-semi-simple double pure imaginary eigenvalues of form $\pm i\omega_0$, where $\omega_0 = 1/\sqrt{2}$. Thus, as $\mu = \mu^*$, condition of type (2.3) holds, or, what is the same, the conditions of Case P_1 , namely, of Subcase P_2^1 , see Subsection 3.2. As $\mu = \mu^*$, system (6.3) has a multiplier $\eta_0 = e^{i\pi\sqrt{2}}$, which is semi-simple, double and indefinite.

Thus, problem on stability of system (6.1) as $\mu = \mu_0$ and as $\mu = \mu^*$ for small values of ε is a problem on parametric resonance. For $\mu = \mu_0$, the resonance is simple, while for $\mu = \mu^*$ the resonance is combinatorial.

We proceed to studying the stability of system (6.1) under the conditions of Cases 1° and 2°.

6.3. Case 1°. We consider the stability of system (6.1) as μ is close to μ_0 and ε is small. Namely, we study this problem for the values (μ, ε) located on the straight line

$$\mu = \mu_0 + m\varepsilon, \tag{6.5}$$

where m is some fixed coefficient.

Substituting (6.5) into (6.1) and making appropriate transformations with (6.2) taken into consideration, we obtain the system

$$\frac{dx}{dt} = J[A_0 + \varepsilon S_1(t) + S_2(t, \varepsilon)]x, \quad x \in \mathbb{R}^4, \tag{6.6}$$

where

$$JA_0 = JA(\mu_0) = \begin{bmatrix} 0 & 1 & 1 & 0 \\ -1 & 0 & 0 & 1 \\ -\frac{1}{2} & \frac{\sqrt{6}}{2} & 0 & 1 \\ \frac{\sqrt{6}}{2} & \frac{5}{4} & -1 & 0 \end{bmatrix},$$

$$JS_1(t) = -mA_1 - \cos t A_2,$$

$$A_1 = \frac{3\sqrt{3}}{2} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{3}{2} & \frac{\sqrt{6}}{2} & 0 & 0 \\ \frac{\sqrt{6}}{2} & \frac{9}{4} & 0 & 0 \end{bmatrix},$$

the matrix $S_2(t, \varepsilon)$ is symmetric, continuous and 2π -periodic in t , smooth in ε and satisfies the relation $\|S_2(t, \varepsilon)\| = O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$ uniformly in t .

System (6.6) is one of form (2.5). In order to study the stability of system (6.6) we employ Theorem 5.1. First we should construct matrix (5.3). This requires to construct an eigenvector $e + ig$ associated with the eigenvalue $i/2$ of the matrix JA_0 . As e and g we can take, for instance, the vectors

$$e = \begin{bmatrix} 8 \\ -2\sqrt{6} \\ 3\sqrt{6}/2 \\ 7 \end{bmatrix}, \quad g = \begin{bmatrix} \sqrt{6} \\ 2 \\ 2 \\ 0 \end{bmatrix}.$$

The numbers T, ω_0, μ_0 (which is redenoted here by η_0) and ν involved in formulae (5.3)-(5.6) read as

$$T = 2\pi, \quad \omega_0 = \frac{1}{2}, \quad \eta_0 = -1, \quad \nu = -1/7.$$

We are in position to calculate the entries of matrix (5.3); after simple calculations we obtain:

$$B = \frac{\pi}{28} \begin{bmatrix} -4\sqrt{6} & 39 - 504\sqrt{2}m \\ 39 + 504\sqrt{2}m & 4\sqrt{6} \end{bmatrix}.$$

In particular, number (5.7) here is as follows: $\Delta = 49\pi^2 (33/16 - 648m^2)$.

By Corollary 5.1 we then obtain that a multiple multiplier $\eta_0 = -1$ of system (6.3) with $\mu = \mu_0$ splits in accordance with formulae (3.6) and (5.8) while passing to system (6.6). In our case these formulae are of the form:

$$\eta_1(\varepsilon) = -1 + \eta_1^{(1)}\varepsilon + O(\varepsilon^{3/2}), \quad \eta_2(\varepsilon) = -1 + \eta_1^{(2)}\varepsilon + O(\varepsilon^{3/2});$$

here

$$\eta_1^{(1)} = \frac{1}{7}\sqrt{\Delta}, \quad \eta_1^{(2)} = -\eta_1^{(1)}.$$

Then by Corollaries 5.2 and 5.3 we obtain that the change of the nature of the stability of system (6.6) occurs as $\Delta = 0$, that is, for the values m defined by the identities $m = \pm k_0$,

where

$$k_0 = \sqrt{\frac{11}{3456}}.$$

At that, systemn (6.6) is unstable if $-k_0 < m < k_0$ and it is stable if $m < -k_0$ or $m > k_0$. This implies that the straight lines

$$\mu = \mu_0 + k_0\varepsilon, \quad \mu = \mu_0 - k_0\varepsilon$$

are the sought tangentials to the curves Γ_1 and Γ_2 at the point $(\mu_0, 0)$.

6.4. Case 2°. We consider the stability of problem (6.1) for μ close to μ^* and for small ε . Namely, we shall study this problem for the values (μ, ε) located on the straight line

$$\mu = \mu^* + m\varepsilon, \tag{6.7}$$

where m is some fixed coefficient.

Substituting (6.7) into (6.1) and making corresponding transformations, we obtain system of form (6.6), in which

$$JA_0 = JA(\mu^*) = \begin{bmatrix} 0 & 1 & 1 & 0 \\ -1 & 0 & 0 & 1 \\ -\frac{1}{4} & \frac{\sqrt{23}}{4} & 0 & 1 \\ \frac{\sqrt{23}}{4} & \frac{5}{4} & -1 & 0 \end{bmatrix},$$

$$JS_1(t) = -mA_1 - \cos tA_2,$$

$$A_1 = \frac{3\sqrt{3}}{2} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \frac{3}{4} & \frac{\sqrt{23}}{4} & 0 & 0 \\ \frac{\sqrt{23}}{4} & \frac{9}{4} & 0 & 0 \end{bmatrix}.$$

In order to study the stability of the obtained problem, we employ Theorem 3.3. We should determine numbers (3.18). In its turn, this requires to construct a pair of non-zero linearly independent vectors $e, g \in \mathbb{C}^{2N}$ such that identities (3.16) holds as $\omega_0 = 1/\sqrt{2}$. These vectors and the eigenvector and the adjoint one, respectively, for the monodromy matrix $V_0 = e^{JA_0T}$. They are associated with the double eigenvalue $e^{i\sqrt{2}\pi}$; namely, the identities hold:

$$V_0e = e^{i\sqrt{2}\pi}e, \quad V_0g = e^{i\sqrt{2}\pi}(g + Te).$$

As e and g we can take, for instance, the vectors

$$e = \begin{bmatrix} 8\sqrt{2} - 2i\sqrt{23} \\ 10i \\ \sqrt{46} - 2i \\ 3\sqrt{2} - 2i\sqrt{23} \end{bmatrix}, \quad g = \frac{1}{3\sqrt{2} - 2i\sqrt{23}} \begin{bmatrix} -4(9\sqrt{23} + 16i\sqrt{2}) \\ 68 - 16i\sqrt{46} \\ -(48 + 24i\sqrt{46}) \\ 0 \end{bmatrix}.$$

The numbers T, ω_0, μ_0 (which is redenoted here by η_0) and ν involved in formulae (3.16) and (3.18) are as follows:

$$T = 2\pi, \quad \omega_0 = \frac{1}{\sqrt{2}}, \quad \eta_0 = e^{i\pi\sqrt{2}}, \quad \nu = \frac{1}{160}.$$

By Corollary 3.6 this implies that a multiple multiplier $\eta_0 = e^{i\pi\sqrt{2}}$ of system (6.3) with $\mu = \mu^*$ splits according to formulae (3.15) and (3.18) while passing to system (6.6). In our case these formulae read as

$$\eta_1(\varepsilon) = \eta_0 + \eta_1^{(1)}\varepsilon^{1/2} + O(\varepsilon), \quad \eta_2(\varepsilon) = \eta_0 + \eta_1^{(2)}\varepsilon^{1/2} + O(\varepsilon);$$

here

$$\eta_1^{(1)} = \frac{1}{\sqrt{2}} e^{i\pi\sqrt{2}} \sqrt[4]{621} \sqrt{m\pi}, \quad \eta_2^{(1)} = -\eta_1^{(1)}.$$

Then by Corollaries 3.7 and 3.8 we obtain that the nature of the stability of system (6.6) changes as $m = 0$. Then system (6.6) is unstable (stable) if $m > 0$ ($m < 0$). This yields that the vertical straight line $\mu = \mu^*$ is the sought tangential to the curve Γ_3 at the point $(\mu^*, 0)$.

7. PROOF OF MAIN STATEMENTS

7.1. Auxiliary constructions. We first recall some auxiliary statements obtained earlier by the authors of the present paper, see [20], [21].

Let $A(\varepsilon)$ be a real square matrix smoothly depending on the parameter ε . First we suppose that the matrix $A_0 = A(0)$ has a semi-simple double eigenvalue λ_0 (real or complex). Then for small $|\varepsilon|$ the matrix $A(\varepsilon)$ possesses two eigenvalues $\lambda^{(1)}(\varepsilon)$ and $\lambda^{(2)}(\varepsilon)$ such that $\lambda^{(1)}(0) = \lambda^{(2)}(0) = \lambda_0$. The mentioned functions are continuously differentiable and are represented as

$$\lambda^{(1)}(\varepsilon) = \lambda_0 + \varepsilon\lambda_1^{(1)} + O(\varepsilon^{3/2}), \quad \lambda^{(2)}(\varepsilon) = \lambda_0 + \varepsilon\lambda_1^{(2)} + O(\varepsilon^{3/2}). \quad (7.1)$$

In the considered case there exist two pairs of linearly independent vectors $e, g \in \mathbb{C}^N$ and $e^*, g^* \in \mathbb{C}^N$ such that

$$A_0 e = \lambda_0 e, \quad A_0 g = \lambda_0 g, \quad A_0^* e^* = \bar{\lambda}_0 e^*, \quad A_0^* g^* = \bar{\lambda}_0 g^*.$$

The vectors e, g, e^*, g^* can be normalized in accordance with the identities

$$(e, e^*) = (g, g^*) = 1, \quad (e, g^*) = (g, e^*) = 0. \quad (7.2)$$

Theorem 7.1. *The coefficients $\lambda_1^{(1)}$ and $\lambda_1^{(2)}$ in formulae (7.1) are the eigenvalues of the matrix*

$$D = \begin{bmatrix} (A_1 e, e^*) & (A_1 g, e^*) \\ (A_1 e, g^*) & (A_1 g, g^*) \end{bmatrix}, \quad (7.3)$$

where $A_1 = A'(0)$.

Now suppose that the matrix $A_0 = A(0)$ possesses a non-semi-simple double eigenvalue λ_0 (real or complex). Then for small $|\varepsilon|$ the matrix $A(\varepsilon)$ has two eigenvalues $\lambda^{(1)}(\varepsilon)$ and $\lambda^{(2)}(\varepsilon)$ such that $\lambda^{(1)}(0) = \lambda^{(2)}(0) = \lambda_0$. These functions are continuous and can be represented by Puiseux expansions:

$$\lambda^{(1)}(\varepsilon) = \lambda_0 + \varepsilon^{1/2}\lambda_1^{(1)} + O(\varepsilon), \quad \lambda^{(2)}(\varepsilon) = \lambda_0 + \varepsilon^{1/2}\lambda_1^{(2)} + O(\varepsilon). \quad (7.4)$$

In the considered case, there exist two pairs of linearly independent vectors $e, g \in \mathbb{C}^N$ and $e^*, g^* \in \mathbb{C}^N$ such that

$$A_0 e = \lambda_0 e, \quad A_0 g = \lambda_0 g + e, \quad A_0^* e^* = \bar{\lambda}_0 e^*, \quad A_0^* g^* = \bar{\lambda}_0 g^* + e^*.$$

The vectors e, g, e^*, g^* can be normalized according to the identities:

$$(e, g^*) = (g, e^*) = 1, \quad (e, e^*) = (g, g^*) = 0. \quad (7.5)$$

Theorem 7.2. *The coefficients $\lambda_1^{(1)}$ and $\lambda_1^{(2)}$ in expansions (7.4) are the numbers*

$$\lambda_1^{(1)} = \sqrt{(A_1 e, e^*)}, \quad \lambda_1^{(2)} = -\lambda_1^{(1)};$$

here $A_1 = A'(0)$.

Without explicit refereeing, we shall make use of the following properties of matrix (1.2):

$$\det J = \det J^{-1} = 1, \quad J^{-1} = J^* = -J, \quad \operatorname{Re}(Jx, x) = 0 \quad \text{for all } x \in \mathbb{C}^{2N}.$$

7.2. Proof of Theorems 3.1 and 3.2. We restrict ourselves by proving Theorem 3.1; Theorem 3.2 can be proved in the same way.

By Theorem 7.1, the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in expansions (3.6) are the eigenvalues of the matrix

$$D = \begin{bmatrix} (V_1 e, e^*) & (V_1 g, e^*) \\ (V_1 e, g^*) & (V_1 g, g^*) \end{bmatrix}; \tag{7.6}$$

here $V_1 = V'(0)$.

For the sake of definiteness, we suppose that under the assumptions of Theorem 3.1 normalization (3.2) holds. Then as the vectors e^* and g^* in matrix (7.6) we employ

$$e^* = -iJe, \quad g^* = -iJg, \tag{7.7}$$

where e and g are the vectors from (3.1). Vectors (7.7) are linearly independent eigenvectors of the matrix $(JA_0)^*$ associated with the eigenvalue $\lambda = -\omega_0 i$, that is, they satisfy the identities

$$(JA_0)^* e^* = -\omega_0 i e^*, \quad (JA_0)^* g^* = -\omega_0 i g^*. \tag{7.8}$$

By identities (3.2) and (3.5), the vectors e, g, e^*, g^* satisfy normalization conditions (7.2).

The monodromy matrix $V(\varepsilon)$ of system (2.5) can be represented as follows, see, for instance, [21]):

$$V(\varepsilon) = V_0 + \varepsilon V_1 + V_2(\varepsilon); \tag{7.9}$$

here $V_0 = e^{JA_0 T}$,

$$V_1 = V'(0) = e^{JA_0 T} \int_0^T e^{-JA_0 \tau} JS_1(\tau) e^{JA_0 \tau} d\tau, \tag{7.10}$$

and $V_2(\varepsilon)$ is a continuously differentiable matrix satisfying the condition: $\|V_2(\varepsilon)\| = O(|\varepsilon|^2)$ as $\varepsilon \rightarrow 0$.

We substitute matrix (7.10) into (7.6) and taking into consideration identities (7.7), we complete the proof of Theorem 3.1.

7.3. Proof of Theorem 3.3. By the vectors e and g in identities (3.16) we define new vectors:

$$e_1 = e, \quad g_1 = \frac{1}{T\mu_0} g, \quad e_1^* = -\alpha Je, \quad g_1^* = \frac{1}{T\mu_0} \alpha Jg,$$

where α is a non-zero coefficient, which is, in general, complex-valued. These vectors satisfy the identities:

$$V_0 e_1 = \mu_0 e_1, \quad V_0 g_1 = \mu_0 g_1 + e_1, \quad V_0^* e_1^* = \overline{\mu_0} e_1^*, \quad V_0^* g_1^* = \overline{\mu_0} g_1^* + e_1^*.$$

At the same time, the vectors e_1, g_1, e_1^*, g_1^* can be normalized in accordance with analogues of identities (7.5) by letting $\alpha = T\overline{\mu_0}/(e, Jg)$.

To complete the proof of Theorem 3.3 it remains to apply Theorem 7.2 to matrix (7.9).

7.4. Proof of Lemma 5.1. By vectors $e, g \in \mathbb{R}^{2N}$ involved in identity (5.1) we define real vectors $e^* = Jg$ and $g^* = Je$. Then

$$(JA_0)^*(g^* + ie^*) = -\omega_0 i(g^* + ie^*),$$

that is, the vector $g^* + ie^*$ is an eigenvectors of the matrix $(JA_0)^*$ associated with the eigenvalue $\lambda = -\omega_0 i$.

According to the spectral theory of linear operators, see, for instance, [19], the space \mathbb{R}^{2N} can be represented as a direct sum $\mathbb{R}^{2N} = E_0 \oplus E^0$ of invariant subspaces of the operator JA_0 . Here E_0 is the root subspace associated with the eigenvalues $\lambda_0 = \pm\omega_0 i$ of the operator JA_0 , while E^0 is the root subspaces associated with the rest of the spectrum of this operator. The

space E_0 is two-dimensional with a basis formed by e and g . This space can be defined by the identity $E^0 = \{v : (v, e^*) = (v, g^*) = 0\}$.

Let us show that $|(e, e^*)| + |(e, g^*)| > 0$. Indeed, assuming the opposite we get $(e, e^*) = (e, g^*) = 0$, that is, $e \in E^0$. On the other hand, by construction $e \in E_0$. This is possible only if $e = 0$, but this contradicts to the fact that e is an eigenvector.

To complete the proof of the lemma, it remains to note that $(e, e^*) = (e, Jg)$ and $(e, g^*) = (e, Je) = 0$.

7.5. Proof of Theorem 5.1. By Theorem 7.1 the coefficients $\mu_1^{(1)}$ and $\mu_1^{(2)}$ in expansions (3.6) are the eigenvalues of matrix (7.6), in which $V_1 = V'(0)$ and the vectors e and g come from identity (5.1). As the vectors e^* and g^* we use the vectors

$$e^* = \nu Jg, \quad g^* = \nu Je, \quad (7.11)$$

where ν is number (5.2).

The entries of matrix (7.6) can be found by means of formulae (7.10) and (7.11). After appropriate transformations we obtain that matrix (7.6) coincides with matrix (5.3).

BIBLIOGRAPHY

1. A.P. Markeev. *Linear Hamiltonian systems and some problems on stability of motion of satellite around the mass center*. Inst. Komput. Issl., Moscow (2009). (in Russian).
2. V.A. Yakubovich, V.M. Starzhinskij. *Linear differential equations with periodic coefficients*. Nauka, Moscow (1972). [John Wiley & Sons, New York; Halsted Press Book, Jerusalem (1975).]
3. V.F. Zhuravlev, F.G. Petrov, M.M. Shunderyuk. *Selected problems in Hamiltonian mechanics*. Lenand, Moscow (2015). (in Russian).
4. V. Lanchares. *On the stability of Hamiltonian dynamical systems // Monografias Matematicas Garca de Galdeano*. **39**, 155–166 (2014).
5. K. Meyer, G. Hall, D. Offin. *Introduction to Hamiltonian dynamical systems and the n-body problem*. Springer, New York (2009).
6. J. Moser. *Lectures on Hamiltonian mechanics*. Nauka, Moscow (1973). (in Russian).
7. V.A. Yakubovich, V.M. Starzhinski. *Parametric resonance in linear systems*. Nauka, Moscow (1987). (in Russian).
8. B.P. Demidovich. *Lectures on mathematical stability theory*. Nauka, Moscow (1967). (in Russian).
9. D. Treschev. *Introduction to the perturbation theory of Hamiltonian systems*. Fazis, Moscow (1998). (in Russian).
10. A.P. Seyranian, A.A. Mailybaev. *Multiparameter stability theory with mechanical applications*. World Scientific, New Jersey (2003).
11. A.P. Markeev. *Libration points in celestial mechanics and cosmodynamics*. Nauka, Moscow (1978). (in Russian).
12. A.P. Markeev. *On a multiple resonance in linear Hamiltonian systems // Dokl. RAN*. **402:3**, 539–543 (2005). [Doklady Phys. **50:5**, 278–282 (2005).]
13. A.D. Bruno. *On types of stability in Hamilton systems // Preprint of Keldysh Inst. Appl. Math.* **021**, 24 pp. (2020). (in Russian).
14. A.D. Bruno. *Normal form of Hamilton system with periodic perturbation // Preprint of Keldysh Inst. Appl. Math.* **057**, 27 pp. (2019). (in Russian).
15. O.V. Kholostova. *On periodic motions of a nonautonomous Hamiltonian system in one case of multiple parametric resonance // Nelinein. Dinam.* **13:4**, 477–504 (2017). (in Russian).
16. B.S. Bardin, E.A. Chekina. *On the constructive algorithm for stability investigation of an equilibrium point of a periodic Hamiltonian system with two degrees of freedom in first-order resonance case // Prikl. Mat. Mekh.* **82:4**, 414–426 (2018). [Mech. Solids. **53**. Suppl. 2. 15–25 (2018).]
17. A. Elipe, V. Lanchares, A. Pascual. *On the stability of equilibria in two-degrees of freedom hamiltonian systems under resonances // J. Nonl. Sci.* **19**, 305–319 (2005).

18. L. Cesari. *Asymptotic behavior and stability problems in ordinary differential equations*. Springer-Verlag, Berlin (1963).
19. T. Kato. *Perturbation theory for linear operators*. Springer-Verlag, Berlin (1966).
20. L.S. Ibragimova L.S. I.G. Fanina, M.G. Yumagulov. *The asymptotic formulae for the problem of the construction hyperbolic and stability regions of dynamical systems* // Ufimskij Matem. Zhurn. **8**:3, 59–81 (2016). [Ufa Math. J. **8**:3, 58–78 (2016).]
21. M.G. Yumagulov, L.S. Ibragimova, A.S. Belova. *Methods for studying the stability of linear periodic systems depending on a small parameter* // Itogi Nauk. Tekhn. Ser. Sovrem. Mat. Pril. Temat. Obz. **163**, 113–126 (2019). (in Russian).
22. A.B. Batkhin, A.D. Bruno, V.P. Varin. *Stability sets of multi-parametric Hamiltonian systems* // Preprints of Keldysh Inst. Appl. Math. **42**, 32 pp. (2011). (in Russian).
23. G.E. Roberts. *Linear stability of the elliptic lagrangian triangle solutions in the three-body problem* // J. Diff. Eqs. **182**:1, 191–218 (2002).
24. T. Kovacs. *Stability chart of the triangular points in the elliptic restricted problem of three bodies* // Mon. Not. R. Astron. Soc. **430**:4, 2755–2760 (2013).
25. M.G. Yumagulov, O.N. Belikova, and N.R. Isanbaeva. *Bifurcation near boundaries of regions of stability of libration points in the three-body problem* // Astron. Zhurn. **95**:2, 158–168 (2018). [Astron. Rep. **62**:2, 144–153 (2018).

Marat Gayazovich Yumagulov,
Bashkir State University,
Zaki Validi str. 32,
450074, Ufa, Russia
E-mail: yum_mg@mail.ru

Lilia Sunagatovna Ibragimova,
Bashkir State University,
Zaki Validi str. 32,
450074, Ufa, Russia
E-mail: lilibr@mail.ru

Anna Sergeevna Belova,
Bashkir State University,
Zaki Validi str. 32,
450074, Ufa, Russia
E-mail: 89177662488@mail.ru